

REVISTA DE LA UNION MATEMATICA ARGENTINA

Director: Darío J. Picco

Vicedirector: Rafael Panzone

Redactores: M. Balanzat, A. Calderón, E. Gentile,
E. Marchi, J. Tirao, C. Trejo

Secretaria de Redacción: M. L. Gastaminza

VOLUMEN 32, NUMERO 4

1986

BAHIA BLANCA

1987

A GENERALIZATION OF THE FUNDAMENTAL FORMULA
 FOR CYLINDERS

Ursula M. Molter

1. ABSTRACT.

Using the G_n -invariant measure on the set of the convex infinite cylinders congruent to Z_q that touch a convex body K , we can generalize the fundamental kinematic formula for cylinders in the sense of Hadwiger in [2].

We obtain a bilinear combination of the "Quermassintegrale" of K and Z_q , with coefficients depending on the first n moments of a non-negative Borel measurable function.

2. INTRODUCTION.

Hadwiger, [2], found a generalization of the kinematic fundamental formula for convex bodies, extending the domain of integration to the whole group of motions in E_n , G_n . He proved that this integral is again a bilinear combination of the "Quermassintegrale" of the convex bodies.

On the other hand, in [1] we find the definition of $n+1$ functionals on convex bodies, which include the "Quermassintegrale" as particular cases.

In this paper we generalize the fundamental kinematic formula for cylinders in the same sense that Hadwiger in [2]. We show that we obtain a bilinear combination of the "Quermassintegrale" of the convex body and the infinite convex cylinder. This generalization includes [1] and [2] as particular cases.

The proof is essentially based on the existence of a tangential measure on the set of infinite convex cylinders congruent to a fixed one Z_q ([3]) and results much easier.

3. NOTATION.

Let E_n be the n -dimensional euclidean space with unit sphere Ω_n . λ_n is the Lebesgue-measure in E_n and $\omega_n = \lambda_n(\Omega_n)$. We denote by K_n the set of all convex, compact and non-empty subsets K in E_n , $K \in K_n$ is called a convex body. For $K \in K_n$, $W_i^n(K)$ are the "Quer-massintegrale" of K ($i = 0, \dots, n$). If $\delta \geq 0$, K_δ is the parallel body in the distance δ of K , and the following Steiner formula holds ([4], p.220-221):

$$(3.1) \quad \begin{aligned} \lambda_n(K_\delta) &= \sum_{j=0}^n \binom{n}{j} W_j^n(K) \delta^j \\ W_i^n(K_\delta) &= \sum_{j=0}^{n-i} \binom{n-i}{j} W_{i+j}^n(K) \delta^j \end{aligned}$$

Now we define infinite convex cylinders as in [4], p.270. Let O be a fixed point in E_n and let L_{n-q} be a $(n-q)$ -plane through O . Let D be a bounded convex body in L_{n-q} . For each point x in D we consider the q -plane orthogonal to L_{n-q} through x . The union of all such L_q is the cylinder Z_q . The q -planes L_q are the generators and D a normal cross section of Z_q . As in [4], p.272, we take

$$(3.2) \quad W_i^n(Z_q) = \begin{cases} W_i^{n-q}(D) & 0 \leq i \leq n-q \\ 0 & n-q < i \leq n \end{cases}$$

We will now explain briefly the fundamental kinematic formula for cylinders ([4], p.272). Let $Z(D)_q$ be the set of all cylinders congruent to Z_q and let γ_D be the normalized G_n -invariant measure on $Z(D)_q$ ([5], p.106). If $K \in K_n$, A is the set of cylinders of $Z(D)_q$ that intersect K , that is $A = \{Z \in Z(D)_q / Z \cap K \neq \emptyset\}$.

We denote with $\gamma_D(K)$, the measure of A in the sense of γ_D , $\gamma_D(K) = \int_A d\gamma_D$, and the fundamental formula for cylinders ([4], p.272) holds:

$$(3.3) \quad \gamma_D(K) = \frac{1}{\omega_n} \sum_{t=0}^{n-q} \binom{n-q}{t} W_{t+q}^n(K) W_{n-t-q}^n(Z_q)$$

4. GENERALIZATION OF THE FUNDAMENTAL FORMULA (3.3).

Let $f: (0, \infty) \rightarrow [0, \infty)$ be an arbitrary Borel measurable function for which holds:

i) $f(0) \neq 0$

ii) The first n moments

$$(4.1) \quad M_k(f) = \int_0^\infty f(r) r^k dr \quad (k = 0, \dots, n-1)$$

are finite.

If we denote with $r = d(K, Z)$ the distance between the convex body K and the cylinder Z , we make the following integral

$$(4.2) \quad R_q(f, K, Z_q) = \frac{\omega_n}{\omega_{n-q}} \int f(r) d\gamma_D,$$

where we integrate over the whole space $Z(D)_q$.

The existence of this integral is assured by the existence of the moments of f (4.1).

PROPOSITION. If $R_q(f, K, Z_q)$ is as in (4.2), the following fundamental formula holds

$$(4.3) \quad R_q(f, K, Z_q) = \frac{1}{\omega_{n-q}} \sum_{t=0}^{n-q} \sum_{s=0}^t C_s(f) \binom{n-q}{t-s} \binom{s+n-t-q}{s} W_{t+q}^n(K) W_{s+n-t-q}^n(Z_q)$$

$$\text{with } C_0(f) = f(0) \quad , \quad C_s(f) = s M_{s-1}(f) \quad (s = 1, \dots, n).$$

That means that $R_q(f, K, Z_q)$ is a bilinear combination of the "Quermassintegrale" of K and Z_q . The moments of f appear in the coefficients.

Proof. We need some previous results. Let $Z(D, K)_q$ be the set of all cylinders congruent to Z_q that touch K (they have non-empty intersection with K , but can be separated weakly by an hyperplane). If A_δ is the set of cylinders congruent to Z_q that intersect K_δ and not K , $A_\delta = \{Z \in Z(D)_q / Z \cap K_\delta \neq \emptyset \text{ and } Z \cap K = \emptyset\}$, then we have $Z(D, K)_q = \lim_{\delta \rightarrow 0} A_\delta$.

In [3] we defined a G_n -invariant measure ϕ_D on $Z(D, K)_q$ which results natural as it verifies $\phi_D(Z(D, K)_q) = \lim_{\delta \rightarrow 0} \frac{1}{\delta} \gamma_D(A_\delta)$.

We also showed in [3] that $\phi_D(Z(D, K)_q)$ is a bilinear combination

$$(4.4) \quad \phi_D(Z(D,K)_q) = \sum_{j=0}^{n-q-1} \binom{n-q}{j+1} \frac{j+1}{\omega_n} W_{n-j}^n(K) W_{j+1}^n(Z_q)$$

Now we may prove (4.3). Following (4.2),

$$R_q(f,K,Z_q) = \frac{\omega_n}{\omega_{n-q}} \int f(r) d\gamma_D.$$

With an obvious reformulation we reach to

$$(4.5) \quad R_q(f,K,Z_q) = \frac{\omega_n}{\omega_{n-q}} [f(0) \gamma_D(K) + \int_0^\infty (f(r) \phi_D(Z(D,K_r)_q) dr]$$

where $Z(D,K_r)_q$ denotes the set of cylinders congruent to Z_q that touch K_r (with K_r the parallel body to K in the distance r).

Using (4.4) we obtain

$$\phi_D(Z(D,K_r)_q) = \sum_{j=0}^{n-q-1} \binom{n-q}{j+1} \frac{j+1}{\omega_n} W_{n-j}^n(K_r) W_{j+1}^n(Z_q)$$

and taking (3.1) into account, it is

$$\phi_D(Z(D,K_r)_q) = \sum_{j=0}^{n-q-1} \sum_{k=0}^j \frac{j+1}{\omega_n} \binom{n-q}{j+1} \binom{j}{k} W_{j+1}^n(Z_q) W_{n-j+k}^n(K) r^k.$$

Now we take $s = k+1$, $t = n+k-j-q$, and get

$$\phi_D(Z(D,K_r)_q) = \sum_{t=1}^{n-q} \sum_{s=1}^t \frac{s r^{s-1}}{\omega_n} \binom{n-q}{t-s} \binom{s+n-t-q}{s} W_{t+q}^n(K) W_{s+n-t-q}^n(Z_q).$$

Hence

$$(4.6) \quad \int_0^\infty f(r) \phi_D(Z(D,K_r)_q) dr = \\ = \sum_{t=1}^{n-q} \sum_{s=1}^t \frac{s M_{s-1}(f)}{\omega_n} \binom{n-q}{t-s} \binom{s+n-t-q}{s} W_{t+q}^n(K) W_{s+n-t-q}^n(Z_q).$$

If we now use formula (3.3) for γ_D and together with (4.6) replace them in (4.5), we reach the desired result.

5. SPECIAL CHOOSSES FOR f , q AND D .

5.1 Special choose for f .

If we take $f(0) = 1$ and $f(r) = 0$ ($r > 0$) then from (4.3) we obtain (3.6) except an irrelevant constant

$$R_q(f,K,Z_q) = \frac{\omega_n}{\omega_{n-q}} \gamma_D(K).$$

5.2 Special choose for q .

In the case $q=0$, that means D is a convex body in E_n , by straightforward calculations we get

$$R_0(f, K, D) = \frac{1}{\omega_n} \sum_{t=0}^n \sum_{s=0}^t C_s(f) \binom{n}{t-s} \binom{s+n-t}{s} W_t^n(K) W_{s+n-t}^n(Z_q).$$

Hence $R_0(f, K, D) = J(f; K, D)$, where $J(f; K, D)$ is the kinematic integral defined by Hadwiger in [2].

5.3 Special choose for D .

Finally, if we take D in L_{n-q} as a point P , Z_q results a q -plane in E_n and taking into account that $W_0^n(Z_q) = W_0^{n-q}(P) = \omega_{n-q}$ and $W_k^n(Z_q) = W_k^{n-q}(P) = 0$ $1 \leq k \leq n$, from (4.3) we obtain

$$R_q(f, K, P) = \sum_{t=0}^n C_t(f) \binom{n-q}{t} W_{t+q}^n(K).$$

That is $R_q(f, K, P) = P_q(f, K)$, where $P_q(f, K)$ are the functionals defined in [1].

REFERENCES

- [1] BOKOWSKY, J., HADWIGER, H und WILLS, J.M., *Eine Erweiterung der Croftonschen Formeln für konvexe Körper*, MATHEMATIKA 23 (1976), 212-219.
- [2] HADWIGER, H., *Eine Erweiterung der kinematischen Hauptformel der Integralgeometrie*, Abhandlungen aus dem Mathematischen Seminar der Univ. Hamburg, 44, Dezember 1975, 84-90.
- [3] MOLTER, U., *Tangential measure on the set of convex infinite cylinders*, Journal of Applied Probab. 23, 961-972 (1986).
- [4] SANTALO, L.A., *Integral Geometry and Geometric Probability*, Reading Mass.: Addison-Wesley (1976).
- [5] SCHNEIDER, R., *Integralgeometrie*, Vorlesungen an der Universität Freiburg im Sommersemester 1979 (1979).