

LIFE SPAN FOR SOLUTIONS OF THE HEAT EQUATION WITH A NONLINEAR BOUNDARY CONDITION

JULIÁN FERNÁNDEZ BONDER AND JULIO D. ROSSI

ABSTRACT. In this note we obtain estimates in terms of the size of the initial data for the blow-up time of positive solutions of the heat equation in \mathbb{R}_+ with a nonlinear boundary condition $-u_x(0, t) = u^p(0, t)$.

INTRODUCTION.

In this note we obtain estimates for the blow-up time of positive solutions of the following parabolic problem

$$(1) \quad \begin{cases} u_t = u_{xx} & \text{in } \mathbb{R}_+ \times (0, T_\lambda), \\ -u_x(0, t) = u^p(0, t) & \text{in } (0, T_\lambda), \\ u(x, 0) = \lambda\phi(x) > 0 & \text{in } \mathbb{R}_+. \end{cases}$$

where $p > 1$ is fixed and $\lambda > 0$ is a parameter.

Throughout this note we assume that the initial datum ϕ is continuous, positive and bounded.

Existence, uniqueness, regularity and continuous dependence on the initial data for this problem were proved, for instance, in [2].

For problem (1), it is well known that if λ is large enough the solution blows up in finite time T_λ (T_λ depends on λ), if and only if $p > 1$, see for example, [1], [3], [4], [8], [10]. This means that there exists a finite time T_λ with

$$\lim_{t \nearrow T_\lambda} \|u(\cdot, t)\|_\infty = +\infty.$$

Here we are interested in the asymptotic behaviour of T_λ when λ goes to infinity. We prove the following Theorem,

Theorem 1. *Under the above assumptions on ϕ , the function $\lambda \mapsto T_\lambda$ is decreasing and continuous with the following asymptotic behaviour at infinity,*

$$\lim_{\lambda \rightarrow \infty} \lambda^{2(p-1)} T_\lambda = T_0.$$

Here T_0 is the blow-up time of the solution of (1) with initial datum $u(x, 0) \equiv \phi(0)$.

Some related papers that deal with the heat equation with a nonlinear source in the entire space are [7] and [9].

1991 *Mathematics Subject Classification.* 35B40, 35K60, 35B30.

Key words and phrases. Parabolic problems, nonlinear boundary conditions, blow-up time.

Partially supported by Universidad de Buenos Aires under grant TX047 and by ANPCyT PICT No. 03-00000-00137. J.D. Rossi is also partially supported by CONICET and by Fundación Antorchas.

Under further assumptions on the initial datum, $u(x, 0) = \psi(x)$ (a compatibility condition and $\psi_{xx} \geq 0$, that guarantee $u_t \geq 0$) it was proved in [6] and [8] that the following blow up rate holds,

$$(2) \quad c \leq (T - t)^{\frac{1}{2(p-1)}} \|u(\cdot, t)\|_\infty \leq C$$

We observe that the exponent that appears in Theorem 1 is related to the one in the blow-up rate (2). This is a consequence of the natural scaling in the equation (1).

PROOF OF THEOREM 1

The fact that $\lambda \mapsto T_\lambda$ is decreasing is an immediate consequence of the maximum principle. To see this, let us call u the solution of (1) with initial datum $\lambda\phi$ and v the solution of (1) with initial datum $\mu\phi$. If $\lambda \leq \mu$ then, by a comparison argument, $u(x, t) \leq v(x, t)$ for all $x > 0$ and $0 < t < \min\{T_\lambda, T_\mu\}$. As T_λ is the blow-up time for u , $\lim_{t \nearrow T_\lambda} \|u(\cdot, t)\|_\infty = +\infty$ and hence v can not be defined beyond T_λ , proving that $T_\mu \leq T_\lambda$.

To see that is continuous we can assume that $\lambda \leq \mu$, hence $T_\lambda \geq T_\mu$. Now, given $\varepsilon > 0$ we have to show that $T_\lambda - \varepsilon < T_\mu$ if $\mu - \lambda < \delta$, but this follows by the continuous dependence with respect to the initial data (see [2]). In fact,

$$\|u(\cdot, T_\lambda - \varepsilon)\|_\infty \leq C = C(\varepsilon)$$

If we replace the power by a globally Lipschitz function $g(u)$ that agrees with u^p for every $u \leq 2C$ we deal with a regular problem, and hence there exists $\delta = \delta(\varepsilon)$ such that

$$\|v(\cdot, T_\lambda - \varepsilon)\|_\infty \leq 2C < +\infty, \quad \text{if } \mu - \lambda < \delta.$$

We observe that as long as $v \leq 2C$ it is a solution of the problem with u^p as nonlinear flux at $x = 0$. By uniqueness, we can conclude that v is bounded up to $T_\lambda - \varepsilon$. Therefore, $T_\mu > T_\lambda - \varepsilon$ as we wanted to prove.

Finally, let us study the asymptotic behaviour at infinity. This is the main point of the paper.

Let u be the solution of (1) and inspired by the natural scaling of the problem we define

$$(3) \quad v_\lambda(x, t) = \frac{1}{\lambda} u(\lambda^{1-p}x, \lambda^{2(1-p)}t).$$

As u satisfies (1), v_λ verifies

$$(4) \quad \begin{cases} (v_\lambda)_t = (v_\lambda)_{xx} & \text{in } \mathbb{R}_+ \times (0, \tilde{T}_\lambda), \\ -(v_\lambda)_x(0, t) = v_\lambda^p(0, t) & \text{in } (0, \tilde{T}_\lambda), \\ v_\lambda(x, 0) = \phi(\lambda^{1-p}x) \equiv \phi_\lambda(x) & \text{in } \mathbb{R}_+. \end{cases}$$

where $\tilde{T}_\lambda = \lambda^{2(p-1)}T_\lambda$.

We want to compute $\lim_{\lambda \rightarrow \infty} \tilde{T}_\lambda$. For that purpose, let us define w as the solution of

$$(5) \quad \begin{cases} w_t = w_{xx} & \text{in } \mathbb{R}_+ \times (0, T_0), \\ -w_x(0, t) = w^p(0, t) & \text{in } (0, T_0), \\ w(x, 0) = \phi(0) & \text{in } \mathbb{R}_+, \end{cases}$$

which is the natural ‘‘limit’’ equation as $\phi_\lambda \rightarrow \phi(0)$ uniformly over compact sets of $[0, +\infty)$.

As $\phi(0) > 0$, w blows up in finite time, T_0 (see [4]).

The Theorem will follow if we prove that

$$\tilde{T}_\lambda \rightarrow T_0, \quad \text{as } \lambda \rightarrow \infty.$$

To prove this claim, let $\varepsilon > 0$ and take $T' = T_0 - \varepsilon$. Let $M = \sup_{0 < t < T'} \|w(\cdot, t)\|_\infty$.

As before, we take $g \in Lip(\mathbb{R})$ such that $g(s) = s^p$ for $s < 2M$. With this g , we define φ the solution of the following problem,

$$(6) \quad \begin{cases} \varphi_t &= \varphi_{xx} & \text{in } \mathbb{R}_+ \times (0, T'), \\ -\varphi_x(0, t) &= g(\varphi)(0, t) & \text{in } (0, T'), \\ \varphi(x, 0) &= \phi_\lambda(x) & \text{in } \mathbb{R}_+. \end{cases}$$

Observe that $\varphi = v_\lambda$ if $v_\lambda < 2M$.

Let us see that $|w(0, t) - \varphi(0, t)| < \delta$ if $\lambda > \lambda_0(\delta)$ for all $t < T'$. For this purpose, let us define $z = w - \varphi$. As $g \in Lip(\mathbb{R})$, z verifies

$$(7) \quad \begin{cases} z_t &= z_{xx} & \text{in } \mathbb{R}_+ \times (0, T'), \\ -z_x(0, t) &= g(w)(0, t) - g(\varphi)(0, t) & \text{in } (0, T'), \\ z(x, 0) &= \phi(0) - \phi_\lambda(x) & \text{in } \mathbb{R}_+. \end{cases}$$

Then we have

$$(8) \quad |z_x(0, t)| \leq K|z(0, t)|,$$

where K depends only on M .

Let $\Gamma(x, t)$ be the fundamental solution of the heat equation, namely

$$\Gamma(x, t) = \frac{1}{(4\pi t)^{1/2}} \exp\left(-\frac{x^2}{4t}\right).$$

For $x \in \mathbb{R}_+$, by (8) we have (see [5])

$$(9) \quad \begin{aligned} z(x, t) &= \int_{\mathbb{R}_+} \Gamma(x-y, t) z(y, 0) dy - \int_0^t \frac{\partial z}{\partial x}(0, \tau) \Gamma(x, t-\tau) d\tau \\ &\quad + \int_0^t \frac{\partial \Gamma}{\partial x}(x, t-\tau) z(0, \tau) d\tau. \end{aligned}$$

Now we observe that Γ satisfies

$$\frac{\partial \Gamma}{\partial x}(0, t-\tau) = 0, \quad \Gamma(0, t-\tau) = \frac{1}{2\sqrt{\pi}(t-\tau)^{\frac{1}{2}}}.$$

Hence, using the initial and boundary conditions we get that

$$|z(0, t)| \leq \int_{\mathbb{R}_+} \Gamma(-y, t) |z(y, 0)| dy + \frac{K}{2\sqrt{\pi}} \int_0^t \frac{|z(0, \tau)|}{(t-\tau)^{\frac{1}{2}}} d\tau.$$

Now we choose $t_0 = t_0(K)$ such that

$$\frac{K}{2\sqrt{\pi}} \int_0^{t_0} \frac{1}{(t_0-\tau)^{\frac{1}{2}}} d\tau \leq \frac{1}{2}.$$

Hence, for $t \in [0, t_0]$ we have

$$\max_{[0, t_0]} |z(0, t)| \leq 2 \max_{[0, t_0]} \int_{\mathbb{R}_+} \Gamma(-y, t) |z(y, 0)| dy$$

We observe that for every $\delta_1 > 0$ there exists $\lambda_1 > 0$ such that

$$\begin{aligned} \int_{\mathbb{R}_+} \Gamma(-y, t) |z(y, 0)| dy &= \int_0^L \Gamma(-y, t) |z(y, 0)| dy + \int_L^{+\infty} \Gamma(-y, t) |z(y, 0)| dy \\ &\leq \eta \int_0^L \Gamma(-y, t) dy + C \int_L^{+\infty} \Gamma(-y, t) dy \\ &\leq \delta_1 \end{aligned}$$

if $\lambda > \lambda_1$.

Now, choose L large so that $\int_L^{+\infty} \Gamma(x - y, t) dy$ is small uniformly in $(x, t) \in (0, L/2) \times (0, t_0)$, and take $\lambda_2 > 0$ such that $|z(y, 0)| < \eta$ for $y \in (0, L)$ and η small.

With this bound on $|z(0, t)|$ we can control $z(x, t)$ for $(x, t) \in (0, L/2) \times (0, t_0)$, in fact, from (8) and (9) we have

$$\begin{aligned} |z(x, t)| &\leq \int_{\mathbb{R}_+} \Gamma(x - y, t) |z(y, 0)| dy + K\delta_1 \int_0^t \Gamma(x, t - \tau) d\tau + \delta_1 \int_0^t \frac{\partial \Gamma}{\partial x}(x, t - \tau) d\tau \\ &\leq \int_0^L \Gamma(x - y, t) |z(y, 0)| dy + \int_L^{+\infty} \Gamma(x - y, t) |z(y, 0)| dy + C\delta_1 \\ &\leq \eta \int_0^L \Gamma(x - y, t) dy + C \int_L^{+\infty} \Gamma(x - y, t) dy + C\delta_1 \leq \delta_2 \end{aligned}$$

if λ is big enough.

Now, as t_0 is independent of λ , we can repeat this procedure beginning with $z(x, t_0)$ as initial datum to find that $|z(x, t)| < \delta_3$ for $(x, t) \in (0, L/4) \times (t_0, 2t_0)$. Therefore, after a finite number of iterations we obtain that, for λ large ($\lambda > \lambda_0(\delta)$)

$$|z(0, t)| < \delta \text{ for all } t < T',$$

as we wanted to see.

Now, as $w(0, T') \leq M$ and $|w(0, t) - \varphi(0, t)| < \delta$, we have that $\varphi(0, t) < 2M$ in $[0, T']$. Therefore, by uniqueness, $\varphi = v_\lambda$ in $[0, T']$. Hence $\tilde{T}_\lambda \geq T' = T_0 - \varepsilon$.

Now, take ψ a compatible initial datum with compact support and $\psi_{xx} \geq 0$ such that $\psi(x) < \phi(0)$ and $\phi(0) - \psi(x)$ small enough in $(0, L)$. From the previous argument, it follows that the solution \underline{w} of (1) with ψ as initial datum verifies

$$w(0, T') - \underline{w}(0, T') < \delta.$$

Hence $T(\psi) \geq T'$. By the assumptions on ψ , \underline{w} verifies (2). Then

$$w(0, T') - \delta \leq \underline{w}(0, T') \leq \|\underline{w}(\cdot, T')\|_\infty \leq C(T(\psi) - T')^{-\frac{1}{2(p-1)}}.$$

Therefore it is easy to see that $T(\psi) - T' < \kappa$ if $\varepsilon = T - T'$ is small (depending on κ). Now, choosing λ large enough, we can obtain $\phi_\lambda(x) > \psi(x)$, then $\tilde{T}_\lambda \leq T(\psi) < T' + \kappa$ and hence as $T_0 - T' = \varepsilon$, we conclude the desired result. \square

REFERENCES

- [1] G. Acosta and J. D. Rossi, *Blow-up vs. global existence for quasilinear parabolic systems with a nonlinear boundary condition*, Z. Angew Math. Phys., Vol 48 (5) (1997), 711-724.
- [2] H. Amann, *Dynamic theory of quasilinear parabolic systems III. Global existence*, Math. Z. Vol. 202(2) (1989), 219-254.
- [3] K. Deng, *Blow-up rates for parabolic systems*, Z. Angew Math. Phys. vol 46. (1995), 110-118.
- [4] K. Deng, M. Fila and H. A. Levine, *On critical exponents for a system of heat equations coupled in the boundary conditions*, Acta Math. Univ Comenianae vol LXIII (1994), 169-192.

- [5] A. Friedman, *Partial differential equations of parabolic type*, Prentice-Hall, Englewood Cliffs, NJ (1964).
- [6] M. Fila and P. Quittner, *The blow-up rate for the heat equation with a nonlinear boundary condition*, *Math. Meth. Appl. Sci.* vol 14. (1991), 197–205.
- [7] C. Gui and X. Wang, *Life span of solutions of the Cauchy problem for a semilinear heat equation*, *Jour. Diff. Eq.*, 115, 166–172 (1995).
- [8] B. Hu and H. M. Yin, *The profile near blow-up time for the solution of the heat equation with a nonlinear boundary condition*, *Trans. Amer. Math. Soc.* vol 346 (1) (1995), 117–135.
- [9] T.-Y. Lee and W.-M. Ni, *Global existence, large time behaviour and life span of solutions of a semilinear parabolic Cauchy problem*, *Trans. Amer. Math. Soc.* 333 (1992), 356–371.
- [10] J. D. Rossi and N. Wolanski, *Global Existence and Nonexistence for a Parabolic System with Nonlinear Boundary Conditions*, *Diff. Int. Eq. Vol.* 11(1), (1998), 179–190.

DEPARTAMENTO DE MATEMÁTICA, FCEYN, UBA (1428) BUENOS AIRES, ARGENTINA.
E-mail address: jfbonder@dm.uba.ar, jrossi@dm.uba.ar