

# THE BEST SOBOLEV TRACE CONSTANT IN DOMAINS WITH HOLES FOR CRITICAL OR SUBCRITICAL EXPONENTS

JULIÁN FERNÁNDEZ BONDER, RAFAEL ORIVE AND JULIO D. ROSSI

ABSTRACT. In this paper we study the best constant in the Sobolev trace embedding  $H^1(\Omega) \hookrightarrow L^q(\partial\Omega)$  in a bounded smooth domain for critical or subcritical  $q$ , that is  $1 < q \leq 2_* = 2(N-1)/(N-2)$ . First, we consider a domain with periodically distributed holes inside where we impose that the involved functions vanish. There exists a critical size of the holes for which the limit problem has an extra term. For sizes larger than critical the best trace constant diverges to infinity and for sizes smaller than critical it converges to the best constant in the domain without holes. Also, we study the problem with the holes located on the boundary of the domain. In this case other critical exists and its extra term appears on the boundary.

## 1. INTRODUCTION.

Sobolev inequalities have been studied by many authors and is by now a classical subject. It at least goes back to [2], for more references see [10]. Relevant for the study of boundary value problems for differential operators is the Sobolev trace inequality that has been intensively studied, see for example, [3, 11, 12, 13] and their references. Given a bounded smooth domain  $\Omega \subset \mathbb{R}^N$ , we deal here with the best constant of the Sobolev trace embedding  $H^1(\Omega) \hookrightarrow L^q(\partial\Omega)$  for critical or subcritical exponents,  $1 < q \leq 2_* := 2(N-1)/(N-2)$ . When  $q = 2$  this leads to an eigenvalue problem of the Steklov type (see [22]). When  $q = 2_*$  this problem is related to the so called Yamabe problem for manifolds with boundaries, see [1, 3, 5, 11].

For subcritical  $q$ ,  $1 < q < 2_*$ , the embedding is compact and hence there exists extremals [14]. When  $q$  is critical,  $q = 2_*$ , the embedding is continuous but no longer compact, hence the existence of extremals is more involved. In [1] it is proved that if the boundary of the domain contains a point with positive mean curvature then there is an extremal for the embedding. Hence, for any bounded smooth domain there exists an extremal even in the critical case,  $q = 2_*$ , see also [13].

Homogenization Theory was created to model and predict the behavior of inhomogeneous materials where inhomogeneities takes places on a small scale. The homogenization of solutions of boundary-value problems in perforated domains have

---

*Key words and phrases.* Homogenization, Nonlinear boundary conditions, Sobolev trace embedding.

2000 *Mathematics Subject Classification.* 35B27, 35J65, 46E35.

The first and third authors are supported by Fundacion Antorchas, CONICET and AN-PCyT PICT 05009 and 10608. The second author was partially supported by the grants S-0505/ESP/0158 of the CAM (Spain) and the MTM2005-00714 and MTM2005-05980 of the MEC.

attracted a lot of attention since the pioneering work [6]. In this paper we consider homogenization problems for the best Sobolev trace constant in perforated domains, following the approach developed in [6].

First, we consider a domain with holes periodically distributed inside the domain. That is, a bounded smooth domain  $\Omega \subset \mathbb{R}^N$ ,  $N \geq 3$ , perturbed periodically with holes located at the interior that decrease its size and increase its number as the homogenization parameter  $\varepsilon$  goes to zero, and find that there exists a critical size of the holes for which the limit problem has an extra term. For sizes larger than critical the best trace constant diverges to infinity and for sizes smaller than critical it converges to the best constant in the domain without holes. When we deal with a subcritical exponent we have compactness of the embedding  $H^1(\Omega) \hookrightarrow L^q(\partial\Omega)$ , however to prove our result for  $q = 2_*$  we need to impose a restriction on the involved domains in order to recover some compactness, in the spirit of [2] (see also [1, 13], etc.).

Next, we prove some homogenization results when the semi-holes are located on the boundary of the domain. In this case the critical size of the holes is different from the critical size for holes inside the domain and moreover for the critical size an extra term appears on the boundary.

**1.1. Holes in the interior.** The Sobolev trace constant in domains with a hole (a subdomain of  $\Omega$  where the functions are forced to vanish) was first studied in [14] where the authors show that there exists an optimal hole that minimizes the best constant among sets with given volume, in the class of measurable sets. On the other hand, it is also proved that a set  $A$  that maximizes  $S$  does not exist. In a subsequent paper, [15], the interior regularity of these optimal holes was studied.

For the study of the behavior of solutions of boundary value problems in domains perturbed periodically with holes complemented with homogeneous boundary conditions (either Dirichlet or Neumann) we refer, for example, to [6, 7, 8].

Let us first describe the domains that we are considering. Let  $B(0, r(\varepsilon)) \subset \mathbb{R}^N$  be the ball of radius  $r(\varepsilon)$ , centered at 0. We assume that  $r(\varepsilon) < \varepsilon$  for any  $\varepsilon \leq 1$ . For each  $\varepsilon$  and for any integer vector  $n \in \mathbb{Z}^N$ , we shall denote by  $B_n^\varepsilon$  the translated image of  $B(0, r(\varepsilon))$  by the vector  $2n\varepsilon$ , i.e.,  $B_n^\varepsilon = B(0, r(\varepsilon)) + 2n\varepsilon$ . Also, let us denote by  $\mathbf{B}^\varepsilon$  the set of all the holes strictly contained in  $\Omega$ , i.e.,

$$\mathbf{B}^\varepsilon = \bigcup \{B_n^\varepsilon \mid \text{dist}(2n\varepsilon, \partial\Omega) \geq \varepsilon, \quad n \in \mathbb{Z}^N\},$$

and we set

$$\Omega^\varepsilon = \Omega \setminus \mathbf{B}^\varepsilon.$$

Hence,  $\Omega^\varepsilon$  is a periodically perforated domain with holes of size  $r(\varepsilon)$ . All of them have the same shape, the distance between two adjacent holes is of order  $\varepsilon$  and they do not overlap. Also, let us remark that the holes are located at a distance at least  $\varepsilon$  from the boundary,  $\partial\Omega$ . When the holes are allowed to touch the boundary the situation is different, see below and Section 3. Let us consider the space of functions,  $H_\varepsilon^1(\Omega) = \{u \in H^1(\Omega) : u|_{\mathbf{B}^\varepsilon} \equiv 0\}$ . The best Sobolev trace constant of

the Sobolev embedding  $H_\varepsilon^1(\Omega) \hookrightarrow L^q(\partial\Omega)$  is then given by

$$(1.1) \quad \lambda(\varepsilon) = \inf_{v \in H_\varepsilon^1(\Omega) \setminus H_0^1(\Omega)} \frac{\int_\Omega |\nabla v|^2 + v^2 dx}{\left( \int_{\partial\Omega} |v|^q dS \right)^{2/q}}.$$

The extremals, normalized by

$$(1.2) \quad \int_{\partial\Omega} |u|^q dS = 1,$$

are positive in  $\Omega^\varepsilon$  and weak solutions to

$$(1.3) \quad \begin{cases} \Delta u_\varepsilon = u_\varepsilon & \text{in } \Omega^\varepsilon, \\ \frac{\partial u_\varepsilon}{\partial \nu} = \lambda(\varepsilon) |u_\varepsilon|^{q-2} u_\varepsilon & \text{on } \partial\Omega, \\ u_\varepsilon = 0 & \text{in } \mathbf{B}^\varepsilon. \end{cases}$$

Our result for interior holes reads:

**Theorem 1.** *Let  $\Omega_\varepsilon$  be a perforated domain with periodic interior holes of radius  $r(\varepsilon) = c_0 \varepsilon^a$ ,  $c_0 > 0$ . Let  $q$  be subcritical,  $1 < q < 2_* := 2(N-1)/(N-2)$ , or critical,  $q = 2_*$ . In the critical case we also assume that*

$$(1.4) \quad (\omega_N(N-2)(c_0/2)^N + 1) \frac{|\Omega|}{|\partial\Omega|^{2/2_*}} < \frac{1}{K(N)},$$

where  $K(N)$  is given by  $K(N) := ((N-2)/2)(\omega_N)^{1/(N-1)}$  and  $\omega_N$  is the volume of the unit sphere in  $\mathbb{R}^N$ . Then,

- (1) *If  $a = N/(N-2)$ , then there exists a constant  $\mu > 0$  (strange term) such that the function  $\lambda(\varepsilon)$  converges as  $\varepsilon \rightarrow 0$  to  $\lambda_\mu$  given by*

$$(1.5) \quad \lambda_\mu = \inf_{v \in H^1(\Omega) \setminus H_0^1(\Omega)} \frac{\int_\Omega |\nabla v|^2 + (1+\mu)v^2 dx}{\left( \int_{\partial\Omega} |v|^q dS \right)^{2/q}},$$

with  $\mu = \omega_N(N-2)c_0^{N-2}/2^N$ . Moreover, the normalized extremals  $u_\varepsilon$  converge weakly along subsequences to a normalized extremal of the limit problem (1.5).

- (2) *If  $a > N/(N-2)$ , the function  $\lambda(\varepsilon)$  converges, as  $\varepsilon \rightarrow 0$ , to  $\lambda_0$ , the best Sobolev trace constant in the domain without holes, that is*

$$(1.6) \quad \lambda_0 = \inf_{v \in H^1(\Omega) \setminus H_0^1(\Omega)} \frac{\int_\Omega |\nabla v|^2 + v^2 dx}{\left( \int_{\partial\Omega} |v|^q dS \right)^{2/q}}.$$

Moreover, the normalized extremals,  $u_\varepsilon$ , converge weakly along subsequences to a normalized extremal of the limit problem (1.6).

- (3) *If  $a < N/(N-2)$ , there holds  $\lambda(\varepsilon) \rightarrow +\infty$  as  $\varepsilon \rightarrow 0$ . Moreover, we get a bound for the speed at which  $\lambda(\varepsilon)$  goes to  $+\infty$ ,*

$$(1.7) \quad \lambda(\varepsilon) \leq C \varepsilon^{a(N-2)-N}.$$

**Remark 1.1.** *The constant  $K(N)$  is the best Sobolev trace constant in the half-space,*

$$(1.8) \quad \frac{1}{K(N)} = \inf_{\nabla w \in L^2(\mathbb{R}_+^N), w \in L^{2^*}(\partial\mathbb{R}_+^N)} \frac{\int_{\mathbb{R}_+^N} |\nabla w|^2 dx}{\left( \int_{\partial\mathbb{R}_+^N} |w|^{2^*} dx' \right)^{2/2^*}}.$$

*The constant  $K(N)$  is computed in [11].*

**Remark 1.2.** *From the proof of Theorem 1, it can be checked that in the critical case, what is needed is that the best Sobolev trace constants,  $\lambda_\varepsilon$ , for the perforated domains,  $\Omega_\varepsilon$ , to be bounded by  $1/K(N)$  uniformly in  $\varepsilon$ , that is,*

$$(1.9) \quad \lim_{\varepsilon \rightarrow 0} \lambda(\varepsilon) < \frac{1}{K(N)}.$$

*Condition (1.4) is the simplest condition that assures (1.9).*

*We note that for every  $\varepsilon > 0$  it holds  $\lambda(\varepsilon) < 1/K(N)$  for every smooth bounded domain of  $\mathbb{R}^N$  by [1]. However, from their arguments, it is not obvious that this inequality can be made uniformly strict.*

**Remark 1.3.** *The extremals of (1.5) are weak solutions of*

$$\begin{cases} \Delta u = (1 + \mu)u & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} = \lambda_\mu |u|^{q-2}u & \text{on } \partial\Omega. \end{cases}$$

**Remark 1.4.** *In the two-dimensional case, i.e.,  $\Omega \subset \mathbb{R}^2$ , we have an analogous result. The critical radius is now*

$$r(\varepsilon) = \exp\left(-\frac{c_0}{\varepsilon^2}\right).$$

*Thus, for radius larger than critical, the best Sobolev constant goes to  $\infty$ , for radius smaller than critical it goes to the best constant of the domain without holes and for the critical radius a strange term,  $\mu$ , appears with  $\mu = \pi/(2c_0)$ . The proof of this fact is completely analogous to the case  $N \geq 3$  with the choice of an appropriate test function, see [6].*

**Remark 1.5.** *Theorem 1 can be generalized to other configuration of the holes. For example, we can consider non-spherical holes, cylinders, trusses, see [6].*

**1.2. Holes on the boundary.** We prove some homogenization results when the holes are located on the boundary of the domain. This problem is related with the study of the behavior of solutions of periodic mixed conditions on the boundary, Dirichlet and Neumann (see for example [9, 20]), and also, with vibration problems of systems with concentrated masses on the boundary (see for example [18, 19]).

To simplify the exposition we consider holes that are placed on a flat part of the boundary. That is, say,  $\Omega \subseteq \{x_N \leq 0\}$ ,  $\Gamma_1 := \partial\Omega \cap \{x_N = 0\}$  is the closure of a (nonempty) smooth open subset of  $\mathbb{R}^{N-1}$ . We consider periodically distributed semi-holes of size  $r(\varepsilon) = c_0\varepsilon^b$  located only on  $\Gamma_1$ . Assume that  $\Gamma_1$  is divided by

a reticula of size  $\varepsilon$ ,  $\Pi_\varepsilon = \{(2m\varepsilon, 0) \in \Gamma_1 \mid \text{with } m \in \mathbb{Z}^{N-1}\}$ , at each point of the reticula,  $x_i \in \Pi_\varepsilon$ , we take a semi-ball of size  $r(\varepsilon)$ ,

$$S(x_i, r(\varepsilon)) = B(x_i, r(\varepsilon)) \cap \{x_N \leq 0\} \subset \mathbb{R}^N,$$

and consider

$$\mathbf{S}^\varepsilon = \bigcup_{x \in \Pi_\varepsilon} S(x_i, r(\varepsilon)) \text{ such that } S(x_i, r(\varepsilon)) \cap (\partial\Omega \setminus \Gamma_1) = \emptyset,$$

and  $\Omega^\varepsilon = \Omega \setminus \mathbf{S}^\varepsilon$ . Thus,  $\Omega^\varepsilon$  is a periodically perforated domain with holes of size  $r(\varepsilon)$ , the distance between two adjacent holes is of order  $\varepsilon$  and they do not overlap with  $\partial\Omega \setminus \Gamma_1$ . The holes are located on the boundary but they are “solid” since the semi-balls are considered in  $\mathbb{R}^N$ . We denote  $\Gamma_1^\varepsilon = \Gamma_1 \cap \mathbf{S}^\varepsilon$ . As above, we consider the space  $H_\varepsilon^1(\Omega) = \{u \in H^1(\Omega) : u|_{\mathbf{S}^\varepsilon} \equiv 0\}$  and the best Sobolev trace constant associated to this space given by (1.1). The extremals normalized by (1.2) are weak solutions of

$$(1.10) \quad \begin{cases} \Delta u_\varepsilon = u_\varepsilon & \text{in } \Omega^\varepsilon, \\ \frac{\partial u_\varepsilon}{\partial \nu} = \lambda(\varepsilon) |u_\varepsilon|^{q-2} u_\varepsilon & \text{on } \partial\Omega \setminus \Gamma_1^\varepsilon, \\ u_\varepsilon = 0 & \text{in } \mathbf{S}^\varepsilon. \end{cases}$$

In this case we prove that the critical size is different from the critical size for holes inside the domain and for the critical size an extra term appears on the homogenized part of the boundary,  $\Gamma_1$ .

**Theorem 2.** *Let  $q$  subcritical,  $1 < q < 2_*$  or critical,  $q = 2_*$ . Let  $\Omega_\varepsilon$  be a perforated domain with periodic boundary holes of radius  $r(\varepsilon) = c_0 \varepsilon^b$ .*

- (1) *If  $b = (N-1)/(N-2)$ , the function  $\lambda(\varepsilon)$  converges as  $\varepsilon \rightarrow 0$  to  $\lambda_{\mu_1}$  the best Sobolev trace constant in the domain with a weight on the boundary,*

$$(1.11) \quad \lambda_{\mu_1} = \inf_{v \in H^1(\Omega) \setminus H_0^1(\Omega)} \frac{\int_\Omega |\nabla v|^2 + v^2 dx + \mu_1 \int_{\Gamma_1} v^2 dS}{\left( \int_{\partial\Omega} |v|^q dS \right)^{2/q}},$$

where  $\mu_1 = \omega_N (N-2) c_0^{N-2} / 2^N$ . Moreover, the normalized extremals  $u_\varepsilon$  converge weakly along subsequences to a normalized extremal of the limit problem (1.11).

- (2) *If  $b > (N-1)/(N-2)$ , the function  $\lambda(\varepsilon)$  converges as  $\varepsilon \rightarrow 0$  to  $\lambda_0$  the best Sobolev trace constant in the domain without holes defined in (1.6). Moreover, the normalized extremals  $u_\varepsilon$  converge weakly along subsequences to a normalized extremal of the limit problem (1.6).*
- (3) *If  $b < (N-1)/(N-2)$ , the function  $\lambda(\varepsilon)$  converges as  $\varepsilon \rightarrow 0$  to  $\lambda_1$  the best Sobolev trace constant in the  $\Omega$  among functions that vanish on  $\Gamma_1$ ,*

$$(1.12) \quad \lambda_{0, \Gamma_1} = \inf_{v \in H^1(\Omega) \setminus H_0^1(\Omega), v|_{\Gamma_1} \equiv 0} \frac{\int_\Omega |\nabla v|^2 + v^2 dx}{\left( \int_{\partial\Omega} |v|^q dS \right)^{2/q}}.$$

Moreover, the normalized extremals  $u_\varepsilon$  converge weakly along subsequences to a normalized extremal of the limit problem (1.12).

**Remark 1.6.** *For the existence of extremals in the problem of critical exponent, we need that  $\lambda(\varepsilon)$  is uniformly bounded by  $1/K(N)$  with respect to  $\varepsilon$ . However, under our geometric hypothesis, we always have a fixed point on  $\partial\Omega - \Gamma_1$  (included in the boundary of  $\Omega_\varepsilon$ ) with positive main curvature and the distance to the holes is uniform (since we place the holes on a flat part of the boundary). So the result of [1] is applied and we have that (1.9) holds. Therefore, in Theorem 2, we do not need to impose any extra condition on the domain  $\Omega$  for the critical case.*

**Remark 1.7.** *The extremals of (1.11) are weak solutions to*

$$\begin{cases} \Delta u = u & \text{in } \Omega, \\ \frac{\partial u}{\partial \nu} + \mu_1 \chi_{\Gamma_1} u = \lambda_{\mu_1} |u|^{q-2} u & \text{on } \partial\Omega, \end{cases}$$

where  $\chi_{\Gamma_1}$  is the characteristic function of  $\Gamma_1$ .

**Remark 1.8.** *If  $\Omega \subset \mathbb{R}^2$ , we have an analogous result. The critical radius is now*

$$r(\varepsilon) = \exp\left(-\frac{c_0}{\varepsilon}\right).$$

Thus, for the critical radius a strange term,  $\mu$ , appears with  $\mu_1 = \pi/(2c_0)$ .

**1.3. Organization of the paper.** The rest of the paper is organized as follows: in Section 2 we deal with perforated domains with interior holes and finally in Section 3 we consider holes on the boundary.

## 2. INTERIOR HOLES

In this section we prove that if we remove from  $\Omega$  a periodic set of holes (where we impose that the considered functions vanish) there exists a limit of the best Sobolev trace constant if the size of the holes is not too large. We consider holes which are balls of radius  $r(\varepsilon) = c_0 \varepsilon^a$  (see Section 1.1).

First, we construct a sequence of appropriate test functions. Next, we prove Theorem 1 distinguishing three cases:  $a = N/(N-2)$ ,  $a > N/(N-2)$  and  $1 \leq a < N/(N-2)$ .

**2.1. Construction of  $w_\varepsilon$ .** In this subsection, following [6], we show that there exists a sequence  $w_\varepsilon$  that verifies the following assumptions:

- (H1)  $w_\varepsilon \rightharpoonup 1$  weakly in  $H^1(\Omega)$ .
- (H2)  $w_\varepsilon \equiv 0$  on the holes  $B_n^\varepsilon$ .
- (H3) There exists a distribution  $\mu \in W^{-1,\infty}(\Omega)$  such that for every sequence  $v_\varepsilon$  that vanishes on the holes and converges weakly in  $H^1(\Omega)$  to a limit  $v$ , one has

$$(2.1) \quad \langle -\Delta w_\varepsilon, \varphi v_\varepsilon \rangle \rightarrow \langle \mu, \varphi v \rangle, \quad \text{for all } \varphi \in C^\infty(\bar{\Omega}),$$

where  $\langle \cdot, \cdot \rangle$  is the duality pairing.

As in [6], the function  $w_\varepsilon$  is defined on each cube  $P_n^\varepsilon = 2n\varepsilon + (-\varepsilon, \varepsilon)^N$ ,  $n \in \mathbb{Z}^N$ , by setting

$$(2.2) \quad \begin{cases} w_\varepsilon = 0 & \text{in } B_n^\varepsilon, \\ \Delta w_\varepsilon = 0 & \text{in } T_n^\varepsilon - B_n^\varepsilon, \\ w_\varepsilon = 1 & \text{in } P_n^\varepsilon - T_n^\varepsilon, \\ w_\varepsilon & \text{continuous at the interfaces,} \end{cases}$$

where  $T_n^\varepsilon = 2n\varepsilon + B(0, \varepsilon)$ , with  $B(0, \varepsilon)$  the ball with radius  $\varepsilon$  and center 0. Now it is easy to compute  $w_\varepsilon$  in polar coordinates in the annulus  $T_n^\varepsilon - B_n^\varepsilon$ . One has

$$(2.3) \quad w_\varepsilon|_{T_n^\varepsilon - B_n^\varepsilon} = \frac{r(\varepsilon)^{-(N-2)} - r^{-(N-2)}}{r(\varepsilon)^{-(N-2)} - \varepsilon^{-(N-2)}} \quad \text{if } N \geq 3,$$

where  $r = |x - 2n\varepsilon|$ . The result in [6] says:

**Lemma 2.1.** *If we choose  $r(\varepsilon) = c_0\varepsilon^{N/N-2}$  for  $N \geq 3$  with  $c_0$  a positive constant, then  $w_\varepsilon$  defined by (2.2) satisfies hypotheses (H1)–(H3) with*

$$(2.4) \quad \mu = \frac{\omega_N(N-2)}{2^N} c_0^{N-2},$$

where  $\omega_N$  is the surface of the unit sphere in  $\mathbb{R}^N$ .

In order to prove Theorem 1, we compute the  $L^2$ -norm of  $\nabla w_\varepsilon$ . As the number  $n(\varepsilon)$  of cubes  $P_n^\varepsilon$  with holes is about  $|\Omega|/(2\varepsilon)^N$ , we get

$$\|\nabla w_\varepsilon\|_{L^2(\Omega)^N}^2 \cong \frac{|\Omega|}{(2\varepsilon)^N} \int_{P_n^\varepsilon} |\nabla w_\varepsilon|^2 dx.$$

Thanks to (2.3) and  $r(\varepsilon) < \varepsilon$ , we obtain

$$\|\nabla w_\varepsilon\|_{L^2(\Omega)^N}^2 \cong \frac{|\Omega|}{(2\varepsilon)^N} \frac{\omega_N(N-2)}{r(\varepsilon)^{-(N-2)}}.$$

Considering that  $r(\varepsilon) = c_0\varepsilon^a$ , we have

$$(2.5) \quad \|\nabla w_\varepsilon\|_{L^2(\Omega)^N}^2 = \begin{cases} \omega_N(N-2)\left(\frac{c_0}{2}\right)^N |\Omega| \varepsilon^{a(N-2)-N} + o(\varepsilon^{a(N-2)-N}), & a \neq \frac{N}{N-2}, \\ \omega_N(N-2)\left(\frac{c_0}{2}\right)^N |\Omega| + o(1), & a = \frac{N}{N-2}. \end{cases}$$

On the other hand, by the definition of  $B^\varepsilon$ , we have that every hole is contained in  $\Omega$  and the distance to the boundary is larger than  $\varepsilon$ . Thus, we get

$$(2.6) \quad w_\varepsilon \equiv 1 \quad \text{and} \quad \frac{\partial w_\varepsilon}{\partial \nu} = 0 \quad \text{on } \partial\Omega.$$

Hence,

$$(2.7) \quad \int_{\partial\Omega} |w_\varepsilon|^q dS = |\partial\Omega|.$$

**2.2. Case  $a = N/(N - 2)$ .** Let us observe that, under the assumptions of Theorem 1, the best Sobolev trace constants of the perforated domains,  $\lambda(\varepsilon)$ , are bounded independently of  $\varepsilon$ . To this end, let us use  $w_\varepsilon$  as a test function in the infimum that defines  $\lambda(\varepsilon)$ , we get, using the estimates on  $w_\varepsilon$  (2.5) and (2.7) proved in the previous subsection,

$$(2.8) \quad \lambda(\varepsilon) \leq \frac{\int_{\Omega} |\nabla w_\varepsilon|^2 + w_\varepsilon^2 dx}{\left( \int_{\partial\Omega} |w_\varepsilon|^q dS \right)^{2/q}} \leq \frac{(\omega_N(N-2)\left(\frac{c_0}{2}\right)^N |\Omega| + 1) |\Omega|}{|\partial\Omega|^{2/q}} + o(1).$$

Hence, the extremals  $u_\varepsilon$ , the weak solutions of (1.3), are bounded in  $H^1(\Omega)$  and we have, for a subsequence,

$$u_\varepsilon \rightharpoonup u \geq 0 \quad \text{weakly in } H^1(\Omega),$$

As the extremals,  $u_\varepsilon$ , are weak solutions of (1.3) and since  $w_\varepsilon$  vanishes on the holes,  $\psi = w_\varepsilon \varphi$  with  $\varphi \in C^\infty(\bar{\Omega})$  is an admissible test function for the weak formulation of (1.3) and satisfies

$$(2.9) \quad \int_{\Omega} \nabla \varphi \nabla u_\varepsilon w_\varepsilon dx + \int_{\Omega} \varphi \nabla u_\varepsilon \nabla w_\varepsilon dx + \int_{\Omega} u_\varepsilon \varphi w_\varepsilon dx = \lambda(\varepsilon) \int_{\partial\Omega} u_\varepsilon^{q-1} \varphi w_\varepsilon dS.$$

Now we observe that, since  $u_\varepsilon \rightarrow u$ ,  $w_\varepsilon \rightarrow 1$  strongly in  $L^2(\partial\Omega)$ , we have

$$\int_{\partial\Omega} u_\varepsilon^{q-1} \varphi w_\varepsilon dS \rightarrow \int_{\partial\Omega} u^{q-1} \varphi dS.$$

Similarly, as  $u_\varepsilon \rightarrow u$ ,  $w_\varepsilon \rightarrow 1$  strongly in  $L^2(\Omega)$ ,

$$\int_{\Omega} u_\varepsilon \varphi w_\varepsilon dx \rightarrow \int_{\Omega} u \varphi dx.$$

Moreover, since  $\nabla u_\varepsilon \rightharpoonup \nabla u$  weakly in  $L^2(\Omega)$  and  $w_\varepsilon \rightarrow 1$  strongly in  $L^2(\Omega)$  we have

$$\int_{\Omega} \nabla \varphi \nabla u_\varepsilon w_\varepsilon dx \rightarrow \int_{\Omega} \nabla \varphi \nabla u dx.$$

To deal with the last term, we integrate by parts using that  $u_\varepsilon$  vanishes on  $\mathbf{B}^\varepsilon$  and obtain

$$\int_{\Omega_\varepsilon} \varphi \nabla u_\varepsilon \nabla w_\varepsilon dx = - \int_{\Omega_\varepsilon} \nabla \varphi \nabla w_\varepsilon u_\varepsilon dx - \int_{\Omega_\varepsilon} \varphi u_\varepsilon \Delta w_\varepsilon dx + \int_{\partial\Omega} \varphi u_\varepsilon \frac{\partial w_\varepsilon}{\partial \nu} dS.$$

We have, by the properties (H1) and (H3) of  $w_\varepsilon$ ,

$$\int_{\Omega} \nabla \varphi \nabla w_\varepsilon u_\varepsilon dx \rightarrow 0, \quad \text{and} \quad - \int_{\Omega} \varphi u_\varepsilon \Delta w_\varepsilon dx \rightarrow \int_{\Omega} \mu u \varphi dx.$$

Also, by (2.6),

$$\int_{\partial\Omega} \varphi u_\varepsilon \frac{\partial w_\varepsilon}{\partial \nu} dS = 0.$$

Now, if we assume that  $\lambda(\varepsilon) \rightarrow \lambda$ , we have

$$(2.10) \quad \int_{\Omega} \nabla \varphi \nabla u dx + \int_{\Omega} \mu u \varphi dx + \int_{\Omega} u \varphi dx = \lambda \int_{\partial\Omega} u^{q-1} \varphi dS.$$

Let us prove that  $\lambda = \lambda_\mu$  defined in (1.5). We now distinguish between the subcritical and the critical cases.



*Subcritical case*,  $1 < q < 2_*$ . In this case, since the immersion  $H^1(\Omega) \hookrightarrow L^q(\partial\Omega)$  is compact and  $\|u_\varepsilon\|_{L^q(\partial\Omega)} = 1$ , we have that  $\|u\|_{L^q(\partial\Omega)} = 1$ . Hence, taking  $\varphi = u$  in (2.10) we get

$$\lambda = \int_{\Omega} |\nabla u|^2 + (1 + \mu)u^2 dx \geq \lambda_\mu.$$

Now, to prove  $\lambda \leq \lambda_\mu$ , let  $u_\mu$  be an extremal of (1.5) and using  $u_\mu w_\varepsilon$  as a test function in (1.1), we get

$$(2.11) \quad \lambda(\varepsilon) \leq \frac{\int_{\Omega} |\nabla(u_\mu w_\varepsilon)|^2 + (u_\mu w_\varepsilon)^2 dx}{\left(\int_{\partial\Omega} |u_\mu w_\varepsilon|^q dS\right)^{2/q}}.$$

By the results of [16] we obtain that  $u_\mu \in C^\alpha(\bar{\Omega})$  and from the maximum principle and Hopf's Lemma we get that  $u_\mu$  is strictly positive in  $\bar{\Omega}$ . Therefore the regularity results of [5] are applicable and we obtain that  $u_\mu \in C^\infty(\bar{\Omega})$ . Thus, by hypothesis (H1), we get

$$\int_{\Omega} (u_\mu w_\varepsilon)^2 dx \rightarrow \int_{\Omega} u_\mu^2 dx \quad \text{and} \quad \int_{\partial\Omega} |u_\mu w_\varepsilon|^q dS \rightarrow \int_{\partial\Omega} |u_\mu|^q dS.$$

On the other hand, we integrate by parts to obtain

$$\begin{aligned} \int_{\Omega} |\nabla(u_\mu w_\varepsilon)|^2 dx &= \int_{\Omega} w_\varepsilon^2 |\nabla u_\mu|^2 dx + 2 \int_{\Omega} u_\mu w_\varepsilon \nabla u_\mu \nabla w_\varepsilon dx \\ &\quad - \int_{\Omega} w_\varepsilon \operatorname{div}(u_\mu^2 \nabla w_\varepsilon) dx + \int_{\partial\Omega} w_\varepsilon u_\mu^2 \frac{\partial w_\varepsilon}{\partial \nu} dS \\ &= \int_{\Omega} w_\varepsilon^2 |\nabla u_\mu|^2 dx - \int_{\Omega} u_\mu^2 w_\varepsilon \Delta w_\varepsilon dx + \int_{\partial\Omega} u_\mu^2 w_\varepsilon \frac{\partial w_\varepsilon}{\partial \nu} dS. \end{aligned}$$

Thus, by the properties (H1) and (H3) of  $w_\varepsilon$  and as  $u_\mu \in C^\infty(\bar{\Omega})$ , we have

$$\int_{\Omega} w_\varepsilon^2 |\nabla u_\mu|^2 dx \rightarrow \int_{\Omega} |\nabla u_\mu|^2 dx \quad \text{and} \quad - \int_{\Omega} u_\mu^2 w_\varepsilon \Delta w_\varepsilon dx \rightarrow \int_{\Omega} \mu u_\mu^2 dx.$$

Also, by (2.6),

$$\int_{\partial\Omega} u_\mu^2 w_\varepsilon \frac{\partial w_\varepsilon}{\partial \nu} dS = 0.$$

Finally, passing to the limit in (2.11), we get  $\lambda \leq \lambda_\mu$  and we conclude the proof of the case  $a = N/(N-2)$  for subcritical  $q$ .

*Critical case*,  $q = 2_*$ . The existence of extremals  $u_\varepsilon$  for (1.1) was proved in [1] (see also [13]). Let us prove that the weak limit of the extremals verifies  $u \neq 0$ . To this end we use the following theorem due to [17].

**Theorem 3.** *There exists a constant  $B > 0$  such that,*

$$\left(\int_{\partial\Omega} v^{2_*} dS\right)^{2/2_*} \leq K(N) \int_{\Omega} |\nabla v|^2 dx + B \int_{\Omega} v^2 dx$$

for every  $v \in H^1(\Omega)$ . Here  $K(N)$  is given by (1.8) and it is sharp.

Now, as  $u_\varepsilon \geq 0$ , it follows that  $u \geq 0$  and, by classical regularity theory,  $u$  is smooth up to the boundary. By the strong maximum principle and Hopf's lemma, it follows that either  $u > 0$  or  $u \equiv 0$ . In order to prove our claim, we have to rule out the possibility of  $u \equiv 0$ . To do this, we adapt the argument given in [13] (see also [2]) to show that  $\|u\|_{L^2(\Omega)} \neq 0$ . In fact, by Theorem 3, as  $u_\varepsilon$  are normalized such that  $\|u_\varepsilon\|_{L^{2^*}(\partial\Omega)} = 1$ , we have

$$1 = \left( \int_{\partial\Omega} u_\varepsilon^{2^*} d\sigma \right)^{2/2^*} \leq K(N) \int_{\Omega} |\nabla u_\varepsilon|^2 dx + B \int_{\Omega} u_\varepsilon^2 dx$$

and hence

$$(2.12) \quad 1 \leq K(N)\lambda(\varepsilon) + (B - K(N)) \int_{\Omega} u_\varepsilon^2 dx.$$

From the estimate (2.8) and the hypothesis (1.4), we get (1.9). Passing to the limit  $\varepsilon \rightarrow 0$  in (2.12) we arrive by (1.9) to

$$(B - K(N)) \int_{\Omega} u^2 dx > 0,$$

and the claim follows.

As before, the limit of the extremals  $u_\varepsilon$ , weak solutions of (1.3), satisfies (we assume, as before, that  $\lambda(\varepsilon) \rightarrow \lambda$ )

$$(2.13) \quad \int_{\Omega} \nabla\varphi \nabla u dx + \int_{\Omega} \mu u \varphi dx + \int_{\Omega} u \varphi dx = \lambda \int_{\partial\Omega} u^{2^*-1} \varphi dS.$$

Taking  $\varphi = u$  in (2.13), we arrive at

$$\int_{\Omega} |\nabla u|^2 + (1 + \mu)u^2 dx = \lambda \int_{\partial\Omega} u^{2^*} dS.$$

As  $u \neq 0$ , we have that  $\lambda > 0$  and  $\|u\|_{L^{2^*}(\partial\Omega)} \neq 0$ . Therefore

$$\lambda_\mu \leq \frac{\int_{\Omega} |\nabla u|^2 + (1 + \mu)u^2 dx}{\left( \int_{\partial\Omega} u^{2^*} dS \right)^{2/2^*}} = \lambda \left( \int_{\partial\Omega} u^{2^*} dS \right)^{1/(N-1)} \leq \lambda.$$

The reverse inequality follows exactly as in the subcritical case.  $\square$

**2.3. Case  $a > N/(N-2)$ .** Again, using as test functions  $w^\varepsilon$ , we get that the best Sobolev trace constants of the perforated domains,  $\lambda(\varepsilon)$ , are bounded independently of  $\varepsilon$ . In fact, by (2.5) and (2.7) we obtain

$$(2.14) \quad \lambda(\varepsilon) \leq \frac{\left( \left( \frac{c_\Omega}{2} \right)^N \varepsilon^{a(N-2)-N} + 1 \right) |\Omega|}{|\partial\Omega|^{2/q}} + o(\varepsilon^{a(N-2)-N}) \leq \frac{2|\Omega|}{|\partial\Omega|^{2/q}} + 1.$$

Thus, the extremals  $u_\varepsilon$  are weak solutions of (1.3) and are bounded in  $H^1(\Omega)$ . We consider a subsequence such that  $\lambda(\varepsilon) \rightarrow \lambda$  and  $u_\varepsilon \rightharpoonup u$  weakly in  $H^1(\Omega)$ .

As in the previous subsection, let us take  $w_\varepsilon \varphi$  with  $\varphi \in C^\infty(\overline{\Omega})$  as a test function and we get the weak formulation (2.9). We pass to the limit in this weak formulation and obtain

$$(2.15) \quad \int_{\Omega} \nabla\varphi \nabla u dx + \int_{\Omega} u \varphi dx = \lambda \int_{\partial\Omega} |u|^{q-2} u \varphi dS,$$

since  $w_\varepsilon$  converges strongly to 1 in  $H^1(\Omega)$  thanks to (2.5) in the case  $a > N/(N-2)$ .

Now, as in the previous subsection, we divide the proof according the exponent  $q$  is subcritical or critical.

*Subcritical case,  $1 < q < 2_*$ .* In this case, as  $\|u_\varepsilon\|_{L^q(\partial\Omega)} = 1$  and as the embedding  $H^1(\Omega) \hookrightarrow L^q(\partial\Omega)$  is compact, we conclude that  $\|u\|_{L^q(\partial\Omega)} = 1$ . Hence, taking  $\varphi = u$  in (2.15), we get that  $\lambda \geq \lambda_0$  by definition of  $\lambda_0$  in (1.6).

To conclude the proof in this case let us prove that  $\lambda \leq \lambda_0$ . Let  $u_0$  be an extremal of (1.6) and let us use  $u_0 w_\varepsilon$  as a test function in (1.1). Thus,

$$\int_{\Omega} |\nabla(u_0 w_\varepsilon)|^2 dx = \int_{\Omega} u_0^2 |\nabla w_\varepsilon|^2 dx + \int_{\Omega} w_\varepsilon^2 |\nabla u_0|^2 dx + 2 \int_{\Omega} u_0 w_\varepsilon \nabla u_0 \nabla w_\varepsilon dx.$$

As  $u_0 \in C^\infty(\bar{\Omega})$  (see [5]), and  $\nabla w_\varepsilon \rightarrow 0$  strongly in  $L^2(\Omega)$  (by (2.5)), we get

$$\int_{\Omega} |\nabla(u_0 w_\varepsilon)|^2 dx \rightarrow \int_{\Omega} |\nabla u_0|^2 dx.$$

Now, we pass to the limit in (2.11) to obtain

$$\lambda \leq \frac{\int_{\Omega} |\nabla u_0|^2 + u_0^2 dx}{\left(\int_{\partial\Omega} |u_0|^q dS\right)^{2/q}} = \lambda_0.$$

This finishes the proof.

*Critical case,  $q = 2_*$ .* In this case, we need to check that  $u \neq 0$ , but this follows as in the previous subsection. In fact, by Theorem 3 we have

$$\begin{aligned} 1 &= \left(\int_{\partial\Omega} |u_\varepsilon|^{2_*} dS\right)^{2/2_*} \leq K(N) \int_{\Omega} |\nabla u_\varepsilon|^2 dx + B \int_{\Omega} |u_\varepsilon|^2 dx \\ &= K(N)\lambda(\varepsilon) + (B - K(N)) \int_{\Omega} |u_\varepsilon|^2 dx \\ &\rightarrow K(N)\lambda + (B - K(N)) \int_{\Omega} |u|^2 dx \end{aligned}$$

and, as from (1.9) it holds that  $K(N)\lambda < 1$ , the claim follows.

Finally, arguing exactly as before, we conclude that  $\lambda = \lambda_0$  and that  $u$  is an extremal for  $\lambda_0$ . This finishes the proof.  $\square$

**2.4. Case  $1 \leq a < N/(N-2)$ .** In this case we have to prove  $\lambda(\varepsilon) \rightarrow \infty$  as  $\varepsilon \rightarrow 0$ . Suppose, contrary to our claim, that there exists a sequence of  $\varepsilon \rightarrow 0$  such that  $\lambda(\varepsilon) \leq C$ . Then, there exists a sequence of normalized functions  $\{u_\varepsilon\}$  in the space  $H_\varepsilon^1(\Omega)$  and satisfying

$$(2.16) \quad \int_{\Omega} |\nabla u_\varepsilon|^2 + u_\varepsilon^2 dx \leq C.$$

Considering  $n(\varepsilon)$ , the number of cubes with holes  $P_n^\varepsilon$  contained in  $\Omega$ , we get

$$\int_{\Omega} |\nabla u_\varepsilon|^2 dx \geq \sum_{i=1}^{n(\varepsilon)} \int_{P_i^\varepsilon} |\nabla u_\varepsilon|^2 dx.$$

Let  $\lambda_1^\varepsilon$  be the Poincaré constant of Sobolev space

$$H := \{u \in H^1(B(0, \varepsilon) - B(0, r(\varepsilon))) \mid u = 0 \text{ on } \partial B(0, r(\varepsilon))\}.$$

It is shown in [21] that

$$(2.17) \quad \lambda_1^\varepsilon \geq C \frac{r(\varepsilon)^{N-2}}{\varepsilon^N} \quad \text{for } N \geq 3.$$

Thus, we obtain

$$\sum_{i=1}^{n(\varepsilon)} \int_{P_i^\varepsilon} |\nabla u_\varepsilon|^2 dx \geq C \frac{r(\varepsilon)^{N-2}}{\varepsilon^N} \sum_{i=1}^{n(\varepsilon)} \int_{P_i^\varepsilon} |u_\varepsilon|^2 dx,$$

and, since  $r(\varepsilon) = \varepsilon^a$  with  $1 \leq a < N/(N-2)$ , we obtain by (2.16) and passing to the limit

$$\lim_{\varepsilon \rightarrow 0} \sum_{i=1}^{n(\varepsilon)} \int_{P_i^\varepsilon} |u_\varepsilon|^2 dx = 0.$$

Therefore,  $u_\varepsilon$  converges to 0 in  $L^2(\Omega)$ . This contradicts the normalization condition (1.2) of the sequence  $u_\varepsilon$ . Hence we obtain that  $\lambda(\varepsilon) \rightarrow \infty$ .

On the other hand, using  $w_\varepsilon$  as a test function in (1.1) we get, using (2.5), that  $w_\varepsilon \rightarrow w$  in  $L^2(\Omega)$  and (2.7),

$$\lambda(\varepsilon) \leq \frac{\int_{\Omega} |\nabla w_\varepsilon|^2 + w_\varepsilon^2 dx}{\left( \int_{\partial\Omega} |w_\varepsilon|^q dS \right)^{2/q}} \leq C \varepsilon^{a(N-2)-N}.$$

Thus, with these estimates we conclude (1.7).  $\square$

### 3. HOLES ON THE BOUNDARY

In this Section we consider holes on the boundary. Recall that we assume that we are dealing with holes on a flat part of the boundary. We distinguish three cases:  $b > (N-1)/(N-2)$ ,  $b = (N-1)/(N-2)$  and  $1 \leq b < (N-1)/(N-2)$ . The proof of the critical exponent  $2_*$  is the same that in the case of interior holes. Here, with our geometric hypothesis of the domain, (1.9) is held (see Remark 1.6).

**3.1. Case  $b > (N-1)/(N-2)$ .** Using the same test function  $w_\varepsilon$  extended by  $w_\varepsilon \equiv 1$  to the whole  $\Omega$  we have

$$\|\nabla w_\varepsilon\|_{L^2(\Omega)^N} \leq \frac{Cr(\varepsilon)^{N-2}}{\varepsilon^{N-1}} = C\varepsilon^{b(N-2)-(N-1)} \rightarrow 0.$$

Hence  $w_\varepsilon \rightarrow 1$  strongly in  $H^1(\Omega)$ . Using  $w_\varepsilon$  as a test function in the definition of  $\lambda_1(\varepsilon)$  we obtain that there exists  $C$  independent of  $\varepsilon$  such that  $\lambda_1(\varepsilon) \leq C$ . From this point the proof follows exactly the same lines as the case  $a > N/(N-2)$  in Section 2.3. However, since the holes are located on the boundary of the domain, (2.6) and (2.7) are not satisfied. We know that  $w_\varepsilon$  converges to 1 in  $L^2(\partial\Omega)$ . Now, we show that

$$(3.1) \quad \frac{\partial w_\varepsilon}{\partial \nu} = 0 \quad \text{on } \partial\Omega.$$

This clearly holds on  $\partial\Omega \setminus \Gamma_1$  by definition of  $w_\varepsilon$ . Now, for any  $\varphi \in L^2(\Gamma_1)$ , we have

$$\int_{\Gamma_1} \varphi \frac{\partial w_\varepsilon}{\partial \nu} dS = \sum_{n=1}^{m(\varepsilon)} \int_{(T_n^\varepsilon - B_n^\varepsilon) \cap \Gamma_1} \varphi \frac{\partial w_\varepsilon}{\partial \nu} dS.$$

where  $m(\varepsilon) \sim \varepsilon^{1-N}$  is the number of holes placed on  $\Gamma_1$ . Let us consider a single cell such that we may assume that the center of the cell is centered at  $x = 0$ . Since  $\Gamma_1$  is considered flat, the normal unit vector is  $-e_N = (0, \dots, 0, -1)$ . Therefore, such like  $w_\varepsilon$  is considered, we get  $\langle -e_N, \nabla w_\varepsilon \rangle = 0$ , and conclude (3.1).

**3.2. Case  $b = (N-1)/(N-2)$ .** As in the case of holes in the interior of  $\Omega$ , the strange term comes from the term that involves  $\Delta w_\varepsilon$ .

We have that  $\Delta w_\varepsilon$  vanishes except on the spheres  $\partial T_i^\varepsilon$  and  $\partial B_i^\varepsilon$ . From the explicit form of  $w_\varepsilon$  we obtain,

$$-\Delta w_\varepsilon = \mu_\varepsilon - \gamma_\varepsilon.$$

The function  $\gamma_\varepsilon$  is supported on the spherical boundary of the semi-ball  $S_i^\varepsilon$ . We observe that, since  $u_\varepsilon = 0$  on this region,

$$\langle \gamma_\varepsilon, u_\varepsilon \rangle = 0.$$

For other side, we know by [6] that in this case

$$\mu_\varepsilon = \sum_{i=1}^{m(\varepsilon)} \frac{\partial w_\varepsilon}{\partial n} \Big|_{\partial T_i^\varepsilon} \delta_i^\varepsilon = \frac{(N-2)c_0^{N-2}}{1-c_0^{N-2}\varepsilon} \sum_{i=1}^{m(\varepsilon)} \delta_i^\varepsilon, \quad \text{in } \mathbb{R}^N,$$

where  $\delta_i^\varepsilon$  are the Dirac masses supported by  $\partial T_i^\varepsilon$  for  $i = 1, \dots, m(\varepsilon)$ . Hence, we have

$$\int_{\Omega} \varphi u_\varepsilon \Delta w_\varepsilon dx = \sum_{i=1}^{m(\varepsilon)} \int_{\partial T_i^\varepsilon \cap \Omega} \frac{(N-2)c_0^{N-2}}{1-c_0^{N-2}\varepsilon} \varphi u_\varepsilon dS.$$

Thanks to the strong convergence of  $u_\varepsilon$  to  $u$  we can pass to the limit and obtain since in [6]:

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega} \varphi u_\varepsilon \Delta w_\varepsilon dx = \frac{\omega_N(N-2)c_0^{N-2}}{2^N} \int_{\Gamma_1} \varphi u dS = \mu_1 \int_{\Gamma_1} \varphi u dS.$$

This case is analogous to spherical holes periodically distributed on a hyperplane of  $\mathbb{R}^N$  of [6]. We note that the capacity of a semi-sphere is a half of the one for the sphere. The rest of the terms can be handled as in the previous section.

**3.3. Case  $1 \leq b < (N-1)/(N-2)$ .** Assume that there exists a sequence  $\varepsilon \rightarrow 0$  with normalized extremals  $\{u_\varepsilon\}$  in the space  $H_\varepsilon^1(\Omega)$ . By definitions of  $\lambda(\varepsilon)$  and  $\lambda_{0,\Gamma_1}$ , we have  $\lambda(\varepsilon) \leq \lambda_{0,\Gamma_1}$ . Therefore,

$$(3.2) \quad \limsup_{\varepsilon \rightarrow 0} \lambda(\varepsilon) \leq \lambda_{0,\Gamma_1},$$

and the extremals satisfy

$$(3.3) \quad \int_{\Omega} |\nabla u_\varepsilon|^2 + u_\varepsilon^2 dx \leq C.$$

Considering  $m(\varepsilon) \sim \varepsilon^{1-N}$ , the number of cells  $P_i^\varepsilon$  placed on  $\Gamma_1$ , we get

$$(3.4) \quad \int_{\Omega} |\nabla u_\varepsilon|^2 dx \geq \sum_{i=1}^{m(\varepsilon)} \int_{P_i^\varepsilon \cap \Omega} |\nabla u_\varepsilon|^2 dx \geq C \frac{\varepsilon^{b(N-2)}}{\varepsilon^N} \sum_{i=1}^{m(\varepsilon)} \int_{P_i^\varepsilon \cap \Omega} |u_\varepsilon|^2 dx,$$

by the Poincaré constant (2.17). Thus, we obtain

$$\sum_{i=1}^{m(\varepsilon)} \int_{P_i^\varepsilon \cap \Omega} |u_\varepsilon|^q dx \geq C \int_{\Gamma_1} \int_0^\varepsilon |u_\varepsilon|^2 dx_N dS_{x'}.$$

Considering the following change of variable  $x_N = \varepsilon y_N$ , we have

$$\int_{\Gamma_1} \int_0^\varepsilon |u_\varepsilon|^q dx_N dx' = \varepsilon \int_{\Gamma_1} \int_0^1 |u_\varepsilon(x', \varepsilon y_N)|^2 dy_N dS_{x'}.$$

Going back to (3.4), we get by (3.3) that

$$C \geq \frac{\varepsilon^{b(N-2)}}{\varepsilon^{N-1}} \int_{\Gamma_1} \int_0^1 |u_\varepsilon(x', \varepsilon y_N)|^2 dy_N dS_{x'}.$$

We pass to the limit as  $\varepsilon \rightarrow 0$ , using that  $1 \leq b < (N-1)/(N-2)$ , we obtain

$$\lim_{\varepsilon \rightarrow 0} \int_{\Gamma_1} \int_0^1 |u_\varepsilon(x', \varepsilon y_N)|^2 dy_N dS_{x'} = 0.$$

Hence, using the regularity of the extremals,

$$\lim_{\varepsilon \rightarrow 0} \int_{\Gamma_1} |u_\varepsilon|^2 dS = 0.$$

Therefore,  $u_\varepsilon$  converges to 0 strongly in  $L^2(\Gamma_1)$ . This shows that every weak limit,  $u$ , of  $u_\varepsilon$  in  $H^1(\Omega)$  verify  $u \equiv 0$  on  $\Gamma_1$ . Therefore

$$(3.5) \quad \lambda_{0,\Gamma_1} \leq \|u\|_{H^1(\Omega)}^2 \leq \liminf_{\varepsilon \rightarrow 0} \|u_\varepsilon\|_{H^1(\Omega)}^2 = \liminf_{\varepsilon \rightarrow 0} \lambda(\varepsilon).$$

From (3.2) and (3.5) we obtain

$$\lim_{\varepsilon \rightarrow 0} \lambda(\varepsilon) = \lambda_{0,\Gamma_1}.$$

Moreover the above arguments show that the limit  $u$  is an extremal of (1.12). This finishes the proof.  $\square$

#### REFERENCES

- [1] Adimurthi, S.L. Yadava. *Positive solution for Neumann problem with critical non linearity on boundary*. Comm. Partial Differential Equations, 16 (11) (1991), 1733–1760.
- [2] T. Aubin. *Équations différentielles non linéaires et le problème de Yamabe concernant la courbure scalaire*. J. Math. Pures et Appl., 55 (1976), 269–296.
- [3] R.J. Biezuner. *Best constants in Sobolev trace inequalities*. Nonlinear Analysis, 54 (2003), 575–589.
- [4] G.A. Chechkin and R. R Gadyl'shin. *On boundary-value problems for the Laplacian in bounded and in unbounded domains with perforated boundaries*. J. Differential Equations 216 (2) (2005), 502–522.
- [5] P. Cherrier, *Problèmes de Neumann non linéaires sur les variétés Riemanniennes*. J. Funct. Anal. Vol. 57 (1984), 154-206.
- [6] D. Cioranescu and F. Murat. *Un terme étrange venu d'ailleurs, I et II*, in Nonlinear Differential Equations and Their Applications, Séminaire Collège de France, H. Brezis and J. L. Lions eds., Vol. II, pp. 98-138, Vol. III, pp. 154-178, Research Notes in Mathematics, 60 and 70, Pitman, London, (1982) and (1983). English translation: *A Strange Term Coming from Nowhere*, in Topics in the Mathematical Modelling of Composite Materials, A. Cherkhaev et. al. eds., Progress in Nonlinear Differential Equations and Their Applications, 31, Birkhäuser, Boston, (1997), 45–93.
- [7] D. Cioranescu and J. Saint Jean Paulin. *Homogenization in Open Sets with Holes*. J. Math. Anal. Appl., 71, (1979), 590–607.

- [8] C. Conca and P. Donato. *Non Homogeneous Neumann Problems in Domains with Small Holes*. RAIRO Modél. Math Anal. Numér., 22, (1988), 561–607.
- [9] A. Damlamian and T.-T. Li. *Boundary homogenization for elliptic problems*. J. Math. Pures Appl. 66 (9) (1987), no. 4, 351–361.
- [10] O. Druet and E. Hebey. *The AB program in geometric analysis: sharp Sobolev inequalities and related problems*. Mem. Amer. Math. Soc. 160 (761) (2002).
- [11] J. F. Escobar. *Sharp constant in a Sobolev trace inequality*. Indiana Math. J., 37 (3) (1988), 687–698.
- [12] J. Fernández Bonder, E. Lami Dozo and J.D. Rossi. *Symmetry properties for the extremals of the Sobolev trace embedding*. Ann. Inst. H. Poincaré. Anal. Non Linéaire, 21 (6) (2004), 795–805.
- [13] J. Fernández Bonder and J.D. Rossi. *On the existence of extremals for the Sobolev trace embedding theorem with critical exponent*. Bull. London Math. Soc., 37 (1) (2005), 119–125.
- [14] J. Fernández Bonder, J.D. Rossi and N. Wolanski. *Behavior of the best Sobolev trace constant and extremals in domains with holes*. To appear in Bull. Sci. Math.
- [15] J. Fernández Bonder, J.D. Rossi and N. Wolanski. *Regularity of the free boundary in an optimization problem related to the best Sobolev trace constant*. SIAM J. Control Optim. 44 (5), 1614–1635, (2005).
- [16] J. Garcia Azorero, I. Peral and J. D. Rossi. *A convex-concave problem with a nonlinear boundary condition*. J. Differential Equations. Vol. 198 (1), 91–128, (2004).
- [17] Y. Li and M. Zhu. *Sharp Sobolev trace inequalities on Riemannian manifolds with boundaries*. Comm. Pure Appl. Math., 50 (1997), 449–487.
- [18] M. Lobo and E. Pérez, *On vibrations of a body with many concentrated masses near the boundary*. Math. Models Methods Appl. Sci. 3 (2) (1993), 249–273.
- [19] M. Lobo and E. Pérez, *Vibrations of a membrane with many concentrated masses near the boundary*. Math. Models Methods Appl. Sci. 5 (5) (1995), 565–585.
- [20] F. Murat, *The Neumann sieve*. Nonlinear variational problems (Isola d’Elba, 1983), 24–32, Res. Notes in Math., 127, Pitman, Boston, MA, 1985.
- [21] J. Rauch and M. Taylor. *Potential and scattering theory on wildly perturbed domains*. J. Funct. Anal., 18, (1975), 27–59.
- [22] M. W. Steklov, *Sur les problèmes fondamentaux en physique mathématique*, Ann. Sci. Ecole Norm. Sup., Vol. 19 (1902), 455–490.

JULIÁN FERNÁNDEZ BONDER

DEPARTAMENTO DE MATEMÁTICA, FCEYN, UNIVERSIDAD DE BUENOS AIRES,  
PABELLON I, CIUDAD UNIVERSITARIA (1428), BUENOS AIRES, ARGENTINA.

*E-mail address:* [jfbonder@dm.uba.ar](mailto:jfbonder@dm.uba.ar)

*Web page:* <http://mate.dm.uba.ar/~jfbonder>

RAFAEL ORIVE

DEPARTAMENTO DE MATEMÁTICAS  
UNIVERSIDAD AUTONOMA DE MADRID  
CRTA. COLMENAR VIEJO KM. 15, 28049 MADRID, SPAIN.

*E-mail address:* [rafael.orive@uam.es](mailto:rafael.orive@uam.es)

*Web page:* <http://www.uam.es/rafael.orive>

JULIO D. ROSSI

INSTITUTO DE MATEMÁTICAS Y FÍSICA FUNDAMENTAL  
CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS  
SERRANO 123, MADRID, SPAIN,  
ON LEAVE FROM DEPARTAMENTO DE MATEMÁTICA, FCEYN UBA (1428)  
BUENOS AIRES, ARGENTINA.

*E-mail address:* [jrossi@dm.uba.ar](mailto:jrossi@dm.uba.ar)

*Web page:* <http://mate.dm.uba.ar/~jrossi>