

# FRACTIONAL SOBOLEV SPACES WITH VARIABLE EXPONENTS AND FRACTIONAL $P(X)$ -LAPLACIANS

URIEL KAUFMANN, JULIO D. ROSSI AND RAUL VIDAL

ABSTRACT. In this article we extend the Sobolev spaces with variable exponents to include the fractional case, and we prove a compact embedding theorem of these spaces into variable exponent Lebesgue spaces. As an application we prove the existence and uniqueness of a solution for a nonlocal problem involving the fractional  $p(x)$ -Laplacian.

## 1. INTRODUCTION

Our main goal in this paper is to extend Sobolev spaces with variable exponents to cover the fractional case.

For a smooth bounded domain  $\Omega \subset \mathbb{R}^n$  we consider two variable exponents, that is, we let  $q : \bar{\Omega} \rightarrow (1, \infty)$  and  $p : \Omega \times \bar{\Omega} \rightarrow (1, \infty)$  be two continuous functions. We assume that  $p$  is symmetric,  $p(x, y) = p(y, x)$ , and that both  $p$  and  $q$  are bounded away from 1 and  $\infty$ , that is, there exist  $1 < q_- < q_+ < +\infty$  and  $1 < p_- < p_+ < +\infty$  such that  $q_- \leq q(x) \leq q_+$  for every  $x \in \bar{\Omega}$  and  $p_- \leq p(x, y) \leq p_+$  for every  $(x, y) \in \bar{\Omega} \times \bar{\Omega}$ .

We define the Banach space  $L^{q(x)}(\Omega)$  as usual,

$$L^{q(x)}(\Omega) := \left\{ f : \Omega \rightarrow \mathbb{R} : \exists \lambda > 0 : \int_{\Omega} \left| \frac{f(x)}{\lambda} \right|^{q(x)} dx < \infty \right\},$$

with its natural norm

$$\|f\|_{L^{q(x)}(\Omega)} := \inf \left\{ \lambda > 0 : \int_{\Omega} \left| \frac{f(x)}{\lambda} \right|^{q(x)} dx < 1 \right\}.$$

Now for  $0 < s < 1$  we introduce the variable exponent Sobolev fractional space as follows:

$$W = W^{s, q(x), p(x, y)}(\Omega) := \left\{ u : \Omega \rightarrow \mathbb{R} : u \in L^{q(x)}(\Omega) : \int_{\Omega} \int_{\Omega} \frac{|f(x) - f(y)|^{p(x, y)}}{\lambda^{p(x, y)} |x - y|^{n + sp(x, y)}} < \infty, \text{ for some } \lambda > 0 \right\},$$

and we set

$$[f]^{s, p(x, y)}(\Omega) := \inf \left\{ \lambda > 0 : \int_{\Omega} \int_{\Omega} \frac{|f(x) - f(y)|^{p(x, y)}}{\lambda^{p(x, y)} |x - y|^{n + sp(x, y)}} < 1 \right\}$$

---

*Key words and phrases.* Variable exponents, Sobolev spaces, fractional Laplacian.  
2010 Mathematics Subject Classification: 46B50, 46E35, 35J60.

as the variable exponent seminorm. It is easy to see that  $W$  is a Banach space with the norm

$$\|f\|_W := \|f\|_{L^{q(x)}(\Omega)} + [f]^{s,p(x,y)}(\Omega);$$

in fact, one just has to follow the arguments in [18] for the constant exponent case. For general theory of classical Sobolev spaces we refer the reader to [1, 4] and for the variable exponent case to [6].

Our main result is the following compact embedding theorem into variable exponent Lebesgue spaces.

**Theorem 1.1.** *Let  $\Omega \subset \mathbb{R}^n$  be a smooth bounded domain and  $s \in (0, 1)$ . Let  $q(x)$ ,  $p(x, y)$  be continuous variable exponents with  $sp(x, y) < n$  for  $(x, y) \in \bar{\Omega} \times \bar{\Omega}$  and  $q(x) > p(x, x)$  for  $x \in \bar{\Omega}$ . Assume that  $r : \bar{\Omega} \rightarrow (1, \infty)$  is a continuous function such that*

$$p^*(x) := \frac{np(x, x)}{n - sp(x, x)} > r(x) \geq r_- > 1,$$

for  $x \in \bar{\Omega}$ . Then, there exists a constant  $C = C(n, s, p, q, r, \Omega)$  such that for every  $f \in W$ , it holds that

$$\|f\|_{L^{r(x)}(\Omega)} \leq C\|f\|_W.$$

That is, the space  $W^{s,q(x),p(x,y)}(\Omega)$  is continuously embedded in  $L^{r(x)}(\Omega)$  for any  $r \in (1, p^*)$ . Moreover, this embedding is compact.

In addition, when one considers functions  $f \in W$  that are compactly supported inside  $\Omega$ , it holds that

$$\|f\|_{L^{r(x)}(\Omega)} \leq C[f]^{s,p(x,y)}(\Omega).$$

**Remark 1.2.** Observe that if  $p$  is a continuous variable exponent in  $\bar{\Omega}$  and we extend  $p$  to  $\bar{\Omega} \times \bar{\Omega}$  as  $p(x, y) := \frac{p(x)+p(y)}{2}$ , then  $p^*(x)$  is the classical Sobolev exponent associated with  $p(x)$ , see [6].

**Remark 1.3.** When  $q(x) \geq r(x)$  for every  $x \in \bar{\Omega}$  the main inequality in the previous theorem,  $\|f\|_{L^{r(x)}(\Omega)} \leq C\|f\|_W$ , trivially holds. Hence our results are meaningful when  $q(x) < r(x)$  for some points  $x$  inside  $\Omega$ .

With the above theorem at hand one can readily deduce existence of solutions to some nonlocal problems. Let us consider the operator  $\mathcal{L}$  given by

$$(1.1) \quad \mathcal{L}u(x) := p.v. \int_{\Omega} \frac{|u(x) - u(y)|^{p(x,y)-2}(u(x) - u(y))}{|x - y|^{n+sp(x,y)}} dy.$$

This operator appears naturally associated with the space  $W$ . In the constant exponent case it is known as the fractional  $p$ -Laplacian, see [2, 3, 5, 7, 8, 9, 11, 12, 15, 16, 17] and references therein. On the other hand, we remark that (1.1) is a fractional version of the well-known  $p(x)$ -Laplacian, given by  $\operatorname{div}(|\nabla u|^{p(x)-2}\nabla u)$ , that is associated with the variable exponent Sobolev space  $W^{1,p(x)}(\Omega)$ . We refer for instance to [6, 10, 13, 14].

Let  $f \in L^{a(x)}(\Omega)$ ,  $a(x) > 1$ . We look for solutions to the problem

$$(1.2) \quad \begin{cases} \mathcal{L}u(x) + |u(x)|^{q(x)-2}u(x) = f(x), & x \in \Omega, \\ u(x) = 0, & x \in \partial\Omega. \end{cases}$$

Associated with this problem we have the following functional (1.3)

$$\mathcal{F}(u) := \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)p(x,y)}} dx dy + \int_{\Omega} \frac{|u(x)|^{q(x)}}{q(x)} dx - \int_{\Omega} f(x)u(x) dx.$$

To take into account the boundary condition in (1.2) we consider the space  $W_0$  that is the closure in  $W$  of compactly supported functions in  $\Omega$ . In order to have a well defined trace on  $\partial\Omega$ , for simplicity, we just restrict ourselves to  $sp_- > 1$ , since then it is easy to see that  $W \subset W^{\bar{s}, p_-}(\Omega) \subset W^{\bar{s}-1/p_-, p_-}(\partial\Omega)$ , with  $\bar{s}p_- > 1$ , see [18, 1]. Concerning problem (1.2), we shall prove the following existence and uniqueness result.

**Theorem 1.4.** *Let  $s \in (1/2, 1)$ , and let  $q(x)$  and  $p(x, y)$  be continuous variable exponents as in Theorem 1.1 with  $sp_- > 1$ . Let  $f \in L^{a(x)}(\Omega)$ , with  $1 < a_- \leq a(x) \leq a_+ < +\infty$  for every  $x \in \bar{\Omega}$ , such that*

$$\frac{np(x, x)}{n - sp(x, x)} > \frac{a(x)}{a(x) - 1} > 1.$$

*Then, there exists a unique minimizer of (1.3) in  $W_0$  that is the unique weak solution to (1.2).*

The rest of the paper is organized as follows: In Section 2 we collect previous results on fractional Sobolev embeddings; in Section 3 we prove our main result, Theorem 1.1, and finally in Section 4 we deal with the elliptic problem (1.2).

## 2. PRELIMINARY RESULTS.

In this section we collect some results that will be used along this paper.

**Theorem 2.1** (Holder's inequality). *Let  $p, q, r : \bar{\Omega} \rightarrow (1, \infty)$  with  $\frac{1}{p} = \frac{1}{q} + \frac{1}{r}$ . If  $f \in L^{r(x)}$  and  $g \in L^{q(x)}$ , then  $fg \in L^{p(x)}$  and*

$$\|fg\|_{L^{p(x)}} \leq c \|f\|_{L^{r(x)}} \|g\|_{L^{q(x)}}.$$

For the constant exponent case we have a fractional Sobolev embedding theorem.

**Theorem 2.2** (Sobolev embedding, [18]). *Let  $s \in (0, 1)$  and  $p \in [1, +\infty)$  such that  $sp < n$ . Then, there exists a positive constant  $C = C(n, p, s)$  such that, for any measurable and compactly supported function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , we have*

$$\|f\|_{L^{p^*}(\mathbb{R}^n)} \leq C \left( \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f(x) - f(y)|^p}{|x - y|^{n+sp}} \right)^{1/p},$$

where

$$p^* = p^*(n, s) = \frac{np}{(n - sp)}$$

is the so-called "fractional critical exponent".

Consequently, the space  $W^{s,p}(\mathbb{R}^n)$  is continuously embedded in  $L^q(\mathbb{R}^n)$  for any  $q \in [p, p^*]$ .

Using the previous result together with an extension property, we also have an embedding theorem in a domain.

**Theorem 2.3** ([18]). *Let  $s \in (0, 1)$  and  $p \in [1, +\infty)$  such that  $sp < n$ . Let  $\Omega \subset \mathbb{R}^n$  be an extension domain for  $W^{s,p}$ . Then there exists a positive constant  $C = C(n, p, s, \Omega)$  such that, for any  $f \in W^{s,p}(\Omega)$ , we have*

$$\|f\|_{L^q(\Omega)} \leq C \|f\|_{W^{s,p}(\Omega)}$$

for any  $q \in [p, p^*]$ ; i.e., the space  $W^{s,p}(\Omega)$  is continuously embedded in  $L^q(\Omega)$  for any  $q \in [p, p^*]$ .

If, in addition,  $\Omega$  is bounded, then the space  $W^{s,p}(\Omega)$  is continuously embedded in  $L^q(\Omega)$  for any  $q \in [1, p^*]$ . Moreover, this embedding is compact for  $q \in [1, p^*)$ .

### 3. FRACTIONAL SOBOLEV SPACES WITH VARIABLE EXPONENTS.

*Proof of Theorem 1.1.* Being  $p$ ,  $q$  and  $r$  continuous, and  $\Omega$  bounded, there exist two positive constants  $k_1$  and  $k_2$  such that

$$(3.1) \quad q(x) - p(x, x) \geq k_1 > 0$$

and

$$(3.2) \quad \frac{np(x, x)}{n - sp(x, x)} - r(x) \geq k_2 > 0,$$

for every  $x \in \overline{\Omega}$ .

Let  $t \in (0, s)$ . Since  $p$ ,  $q$  and  $r$  are continuous, using (3.1) and (3.2) we can find a constant  $\epsilon = \epsilon(p, r, q, k_2, k_1, t)$  and a finite family of disjoint sets  $B_i$  such that

$$\Omega = \cup_{i=1}^N B_i \quad \text{and} \quad \text{diam}(B_i) < \epsilon,$$

that verify that

$$(3.3) \quad \begin{aligned} \frac{np(z, y)}{n - tp(z, y)} - r(x) &\geq \frac{k_2}{2}, \\ q(x) &\geq p(z, y) + \frac{k_1}{2}, \end{aligned}$$

for every  $x \in B_i$  and  $(z, y) \in B_i \times B_i$ .

Let

$$p_i := \inf_{(z, y) \in B_i \times B_i} (p(z, y) - \delta).$$

From (3.3) and the continuity of the involved exponents we can choose  $\delta = \delta(k_2)$ , with  $p_- - 1 > \delta > 0$ , such that

$$(3.4) \quad \frac{np_i}{n - tp_i} \geq \frac{k_2}{3} + r(x)$$

for each  $x \in B_i$ .

It holds that

- (1) if we let  $p_i^* = \frac{np_i}{n - tp_i}$ , then  $p_i^* \geq \frac{k_2}{3} + r(x)$  for every  $x \in B_i$ ,
- (2)  $q(x) \geq p_i + \frac{k_1}{2}$  for every  $x \in B_i$ .

Hence we can apply Theorem 2.3 for constant exponents to obtain the existence of a constant  $C = C(n, p_i, t, \epsilon, B_i)$  such that

$$(3.5) \quad \|f\|_{L^{p_i^*}(B_i)} \leq C \left( \|f\|_{L^{p_i}(B_i)} + [f]^{t, p_i}(B_i) \right).$$

Now we want to show that the following three statements hold.

(A) There exists a constant  $c_1$  such that

$$\sum_{i=0}^N \|f\|_{L^{p_i^*}(B_i)} \geq c_1 \|f\|_{L^{r(x)}(\Omega)}.$$

(B) There exists a constant  $c_2$  such that

$$c_2 \|f\|_{L^{q(x)}(\Omega)} \geq \sum_{i=0}^N \|f\|_{L^{p_i}(B_i)}.$$

(C) There exists a constant  $c_3$  such that

$$c_3 [f]^{s,p(x,y)}(\Omega) \geq \sum_{i=0}^N [f]^{t,p_i}(B_i).$$

These three inequalities and (3.5) imply that

$$\begin{aligned} \|f\|_{L^{r(x)}(\Omega)} &\leq C \sum_{i=0}^N \|f\|_{L^{p_i^*}(B_i)} \\ &\leq C \sum_{i=0}^N \left( \|f\|_{L^{p_i}(B_i)} + [f]^{t,p_i}(B_i) \right) \\ &\leq C \left( \|f\|_{L^{q(x)}(\Omega)} + [f]^{s,p(x,y)}(\Omega) \right) \\ &= C \|f\|_W, \end{aligned}$$

as we wanted to show.

Let us start with (A). We have

$$|f(x)| = \sum_{i=0}^N |f(x)| \chi_{B_i}.$$

Hence

$$(3.6) \quad \|f\|_{L^{r(x)}(\Omega)} \leq \sum_{i=0}^N \|f\|_{L^{r(x)}(B_i)},$$

and by item (1), for each  $i$ ,  $p_i^* > r(x)$  if  $x \in B_i$ . Then we take  $a_i(x)$  such that

$$\frac{1}{r(x)} = \frac{1}{p_i^*} + \frac{1}{a(x)}.$$

Using Theorem 2.1 we obtain

$$\begin{aligned} \|f\|_{L^{r(x)}(B_i)} &\leq c \|f\|_{L^{p_i^*(x)}(B_i)} \|1\|_{L^{a_i(x)}(B_i)} \\ &= C \|f\|_{L^{p_i^*(x)}(B_i)}. \end{aligned}$$

Thus, recalling (3.6) we get [A].

To show (B) we argue in a similar way using that  $q(x) > p_i$  for  $x \in B_i$ .

In order to prove (C) let us set

$$F(x, y) := \frac{|f(x) - f(y)|}{|x - y|^s},$$

and observe that

$$\begin{aligned}
[f]^{t,p_i}(B_i) &= \left( \int_{B_i} \int_{B_i} \frac{|f(x) - f(y)|^{p_i}}{|x - y|^{n+tp_i+sp_i-sp_i}} dx dy \right)^{\frac{1}{p_i}} \\
&= \left( \int_{B_i} \int_{B_i} \left( \frac{|f(x) - f(y)|}{|x - y|^s} \right)^{p_i} \frac{dx dy}{|x - y|^{n+(t-s)p_i}} \right)^{\frac{1}{p_i}} \\
(3.7) \quad &= \|F\|_{L^{p_i}(\mu, B_i \times B_i)} \\
&\leq C \|F\|_{L^{p(x,y)}(\mu, B_i \times B_i)} \|1\|_{L^{b_i(x,y)}(\mu, B_i \times B_i)} \\
&= C \|F\|_{L^{p(x,y)}(\mu, B_i \times B_i)},
\end{aligned}$$

where we have used Theorem 2.1 with

$$\frac{1}{p_i} = \frac{1}{p(x,y)} + \frac{1}{b_i(x,y)},$$

but considering the measure in  $B_i \times B_i$  given by

$$d\mu(x,y) = \frac{dx dy}{|x - y|^{n+(t-s)p_i}}.$$

Now our aim is to show that

$$(3.8) \quad \|F\|_{L^{p(x,y)}(\mu, B_i \times B_i)} \leq C [f]^{s,p(x,y)}(B_i)$$

for every  $i$ . If this is true, then we immediately derive (C) from (3.7).

Let  $\lambda > 0$  be such that

$$\int_{B_i} \int_{B_i} \frac{|f(x) - f(y)|^{p(x,y)}}{\lambda^{p(x,y)} |x - y|^{n+sp(x,y)}} dx dy < 1.$$

Choose

$$k := \sup \left\{ 1, \sup_{(x,y) \in \Omega \times \Omega} |x - y|^{s-t} \right\} \quad \text{and} \quad \tilde{\lambda} := \lambda k.$$

Then

$$\begin{aligned}
&\int_{B_i} \int_{B_i} \left( \frac{|f(x) - f(y)|}{(\tilde{\lambda} |x - y|^s)} \right)^{p(x,y)} \frac{dx dy}{|x - y|^{n+(t-s)p_i}} \\
&= \int_{B_i} \int_{B_i} \frac{|x - y|^{(s-t)p_i}}{k^{p(x,y)}} \frac{|f(x) - f(y)|^{p(x,y)}}{\lambda^{p(x,y)} |x - y|^{n+sp(x,y)}} dx dy \\
&\leq \int_{B_i} \int_{B_i} \frac{|f(x) - f(y)|^{p(x,y)}}{\lambda^{p(x,y)} |x - y|^{n+sp(x,y)}} dx dy < 1.
\end{aligned}$$

Therefore

$$\|F\|_{L^{p(x,y)}(\mu, B_i \times B_i)} \leq \lambda k,$$

which implies the inequality (3.8).

On the other hand, when we consider functions that are compactly supported inside  $\Omega$  we can get rid of the term  $\|f\|_{L^{q(x)}(\Omega)}$  and it holds that

$$\|f\|_{L^{q(x)}(\Omega)} \leq C [f]^{s,p(x,y)}(\Omega).$$

Finally, we recall that the previous embedding is compact since in the constant exponent case we have that for subcritical exponents the embedding is compact.

Hence, for a bounded sequence in  $W$ ,  $f_i$ , we can mimic the previous proof obtaining that for each  $B_i$  we can extract a convergent subsequence in  $L^{r(x)}(B_i)$ .  $\square$

**Remark 3.1.** Our result is sharp in the following sense: if

$$p^*(x_0) := \frac{np(x_0, x_0)}{n - sp(x_0, x_0)} < r(x_0)$$

for some  $x_0 \in \Omega$ , then the embedding of  $W$  in  $L^{r(x)}(\Omega)$  cannot hold for every  $q(x)$ . In fact, from our continuity conditions on  $p$  and  $r$  there is a small ball  $B_\delta(x_0)$  such that

$$\max_{\overline{B_\delta(x_0)} \times \overline{B_\delta(x_0)}} \frac{np(x, y)}{n - sp(x, y)} < \min_{\overline{B_\delta(x_0)}} r(x).$$

Now, fix  $q < \min_{\overline{B_\delta(x_0)}} r(x)$  (note that for  $q(x) \geq r(x)$  we trivially have that  $W$  is embedded in  $L^{r(x)}(\Omega)$ ). In this situation, with the same arguments that hold for the constant exponent case, one can find a sequence  $f_k$  supported inside  $B_\delta(x_0)$  such that  $\|f_k\|_W \leq C$  and  $\|f_k\|_{L^{q(x)}(B_\delta(x_0))} \rightarrow +\infty$ . In fact, just consider a smooth, compactly supported function  $g$  and take  $f_k(x) = k^a g(kx)$  with  $a$  such that  $ap(x, y) - n + sp(x, y) \leq 0$  and  $ar(x) - n > 0$  for  $x, y \in \overline{B_\delta(x_0)}$ .

Finally, we mention that the critical case

$$p^*(x) := \frac{np(x, x)}{n - sp(x, x)} \geq r(x)$$

with equality for some  $x_0 \in \Omega$  is left open.

#### 4. EQUATIONS WITH THE FRACTIONAL $p(x)$ -LAPLACIAN.

In this section we apply our previous results to solve the following problem. Let us consider the operator  $\mathcal{L}$  given by

$$\mathcal{L}u(x) := p.v. \int_{\Omega} \frac{|u(x) - u(y)|^{p(x,y)-2}(u(x) - u(y))}{|x - y|^{n+sp(x,y)}} dy.$$

Let  $\Omega$  be a bounded smooth domain in  $\mathbb{R}^n$  and  $f \in L^{a(x)}(\Omega)$  with  $a_+ > a(x) > a_- > 1$  for each  $x \in \overline{\Omega}$ . We look for solutions to the problem

$$(4.1) \quad \begin{cases} \mathcal{L}u(x) + |u(x)|^{q(x)-2}u(x) = f(x), & x \in \Omega, \\ u(x) = 0, & x \in \partial\Omega. \end{cases}$$

To this end we consider the following functional

$$(4.2) \quad \mathcal{F}(u) := \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)} p(x, y)} dx dy + \int_{\Omega} \frac{|u(x)|^{q(x)}}{q(x)} dx - \int_{\Omega} f(x)u(x) dx.$$

Let us first state the definition of a weak solution to our problem (4.1). Note that here we are using that  $p$  is symmetric, that is, we have  $p(x, y) = p(y, x)$ .

**Definition 4.1.** We call  $u$  a weak solution to (4.1) if  $u \in W_0^{s, q(x), p(x, y)}(\Omega)$  and

$$(4.3) \quad \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{p(x,y)-2}(u(x) - u(y))(v(x) - v(y))}{|x - y|^{n+sp(x,y)}} dx dy + \int_{\Omega} |u|^{q(x)-2}u(x)v(x) dx = \int_{\Omega} f(x)v(x) dx,$$

for every  $v \in W_0^{s,q(x),p(x,y)}(\Omega)$ .

Now our aim is to show that  $\mathcal{F}$  has a unique minimizer in  $W_0^{s,q(x),p(x,y)}(\Omega)$ . This minimizer shall provide the unique weak solution to the problem (4.1).

*Proof of Theorem 1.4.* We just observe that we can apply the direct method of calculus of variations. Note that the functional  $\mathcal{F}$  given in (4.2) is bounded below and strictly convex (this holds since for any  $x$  and  $y$  the function  $t \mapsto t^{p(x,y)}$  is strictly convex).

From our previous results,  $W_0^{s,q(x),p(x,y)}(\Omega)$  is compactly embedded in  $L^{r(x)}(\Omega)$  for  $r(x) < p^*(x)$ , see Theorem 1.1. In particular, we have that  $W_0^{s,q(x),p(x,y)}(\Omega)$  is compactly embedded in  $L^{\frac{\alpha(x)}{\alpha(x)-1}}(\Omega)$ .

Let us see that  $\mathcal{F}$  is coercive. We have

$$\begin{aligned} \mathcal{F}(u) &= \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)p(x,y)}} dx dy + \int_{\Omega} \frac{|u(x)|^{q(x)}}{q(x)} dx - \int_{\Omega} f(x)u(x) dx \\ &\geq \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)p(x,y)}} dx dy + \int_{\Omega} \frac{|u(x)|^{q(x)}}{q(x)} dx \\ &\quad - \|f\|_{L^{\alpha(x)}(\Omega)} \|u\|_{L^{\frac{\alpha(x)}{\alpha(x)-1}}(\Omega)} \\ &\geq \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)p(x,y)}} dx dy + \int_{\Omega} \frac{|u(x)|^{q(x)}}{q(x)} dx - C\|u\|_W. \end{aligned}$$

Now, let us assume that  $\|u\|_W > 1$ . Then we have

$$\begin{aligned} \frac{\mathcal{F}(u)}{\|u\|_W} &\geq \frac{1}{\|u\|_W} \left( \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{p(x,y)}}{|x - y|^{n+sp(x,y)p(x,y)}} dx dy + \int_{\Omega} \frac{|u(x)|^{q(x)}}{q(x)} dx \right) - C \\ &\geq \|u\|_W^{\min\{p_-, q_-\} - 1} - C. \end{aligned}$$

We next choose a sequence  $u_j$  such that  $\|u_j\|_W \rightarrow \infty$  as  $j \rightarrow \infty$ . Then we have

$$\mathcal{F}(u_j) \geq \|u_j\|_W^{\min\{p_-, q_-\}} - C\|u_j\|_W \rightarrow \infty,$$

and we conclude that  $\mathcal{F}$  is coercive. Therefore, there is a unique minimizer of  $\mathcal{F}$ .

Finally, let us check that when  $u$  is a minimizer to (4.2) then it is a weak solution to (4.1). Given  $v \in W_0^{s,q(x),p(x,y)}(\Omega)$  we compute

$$\begin{aligned} 0 &= \frac{d}{dt} \mathcal{F}(u + tv) \Big|_{t=0} = \int_{\Omega} \int_{\Omega} \frac{d}{dt} \frac{|u(x) - u(y) + t(v(x) - v(y))|^{p(x,y)}}{p(x,y)|x - y|^{n+sp(x,y)}} dx dy \Big|_{t=0} \\ &\quad + \int_{\Omega} \frac{d}{dt} \frac{|u(x) + tv(x)|^{q(x)}}{q(x)} dx \Big|_{t=0} - \int_{\Omega} \frac{d}{dt} f(x)(u(x) + tv(x)) dx \Big|_{t=0} \\ &= \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{p(x,y)-2} (u(x) - u(y))(v(x) - v(y))}{|x - y|^{n+sp(x,y)}} dx dy \\ &\quad + \int_{\Omega} |u(x)|^{q(x)-2} u(x)v(x) dx - \int_{\Omega} f(x)v(x) dx, \end{aligned}$$

as  $u$  is a minimizer of (4.2). Thus, we deduce that  $u$  is a weak solution to the problem (4.1).

The proof of the converse (that every weak solution is a minimizer of  $\mathcal{F}$ ) is standard and we leave the details to the reader.  $\square$

## REFERENCES

- [1] R. Adams, J. Fournier, *Sobolev spaces*. Second edition. Pure and Applied Mathematics (Amsterdam), 140. Elsevier/Academic Press, Amsterdam, 2003. xiv+305 pp.
- [2] L. Brasco, E. Parini, M. Squassina. *Stability of variational eigenvalues for the fractional  $p$ -Laplacian*. Discrete Contin. Dyn. Syst. **36** (2016), 1813–1845.
- [3] L. M. Del Pezzo, A. M. Salort, *The first non-zero Neumann  $p$ -fractional eigenvalue*, Nonlinear Anal. **118** (2015), 130–143.
- [4] F. Demengel, G. Demengel, *Functional spaces for the theory of elliptic partial differential equations*, Universitext, Springer, London, 2012. Translated from the 2007 French original by Reinie Erné.
- [5] A. Di Castro, T. Kuusi, G. Palatucci, *Local behavior of fractional  $p$ -minimizers*, Ann. Inst. H. Poincaré Anal. Non Linéaire **33** (2016), 1279–1299.
- [6] L. Diening, P. Harjulehto, P. Hasto, M. Ruzicka, Lebesgue and Sobolev spaces with variable exponents, Lecture Notes in Mathematics, Vol. 2017, Springer-Verlag, Heidelberg, 2011.
- [7] G. Franzina, G. Palatucci, *Fractional  $p$ -eigenvalues*. Riv. Math. Univ. Parma (N.S.) **5** (2014), 373–386.
- [8] A. Iannizzotto, S. Liu, Shibo, K. Perera, M. Squassina, *Existence results for fractional  $p$ -Laplacian problems via Morse theory*, Adv. Calc. Var. **9** (2016), 101–125.
- [9] A. Iannizzotto, S. Mosconi, M. Squassina, *Global Hölder regularity for the fractional  $p$ -Laplacian*, preprint, arXiv:1411.2956.
- [10] P. Harjulehto, P. Hasto, U. Le, M. Nuortio, *Overview of differential equations with non-standard growth*, Nonlinear Anal. **72** (2010), 4551–4574.
- [11] H. Jylha, *An optimal transportation problem related to the limits of solutions of local and nonlocal  $p$ -Laplace-type problems*, Rev. Mat. Complutense **28** (2015), 85–121.
- [12] E. Lindgren, P. Lindqvist, *Fractional eigenvalues*, Calc. Var. Partial Differential Equations **49** (2014), 795–826.
- [13] J.J. Manfredi, J.D. Rossi, J.M. Urbano.  *$p(x)$ -Harmonic functions with unbounded exponent in a subdomain*, Ann. Inst. Henri Poincaré, C. ANL., **26** (2009), 2581–2595.
- [14] J.J. Manfredi, J.D. Rossi, J.M. Urbano, *Limits as  $p(x) \rightarrow \infty$  of  $p(x)$ -harmonic functions*, Nonlinear Anal. **72** (2010), 309–315.
- [15] G. Molica Bisci, *Sequence of weak solutions for fractional equations*, Math. Research Lett. **21** (2014), 241–253.
- [16] G. Molica Bisci, B.A. Pansera, *Three weak solutions for nonlocal fractional equations*, Adv. Nonlinear Stud. **14** (2014), 619–629.
- [17] G. Molica Bisci, *Fractional equations with bounded primitive*, Appl. Math. Lett. **27** (2014), 53–58.
- [18] E. Di Nezza, G. Palatucci, E. Valdinoci, *Hitchhiker’s guide to the fractional Sobolev spaces*, Bull. Sci. Math. **136** (2012), 521–573.

(J. D. Rossi), DEPTO. DE MATEMÁTICA, FCEYN, UNIVERSIDAD DE BUENOS AIRES, CIUDAD UNIVERSITARIA, PAB 1 (1428), BUENOS AIRES, ARGENTINA, EMAIL: JROSSI@DM.UBA.AR

(U. Kaufmann and R. Vidal), FAMAF, UNIVERSIDAD NACIONAL DE CORDOBA, (5000), CORDOBA, ARGENTINA., EMAIL: KAUFMANN@MATE.UNCOR.EDU, VIDAL@MATE.UNCOR.EDU